

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number  
**WO 01/46698 A2**

(51) International Patent Classification<sup>7</sup>: **G01N 33/68**

(21) International Application Number: **PCT/US00/34503**

(22) International Filing Date:  
18 December 2000 (18.12.2000)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/172,979 20 December 1999 (20.12.1999) US  
60/186,626 3 March 2000 (03.03.2000) US  
09/721,908 24 November 2000 (24.11.2000) US

(71) Applicant (for all designated States except US): **CHEMO-CENTRYX, INC.** [US/US]; 1539 Industrial Road, San Carlos, CA 94070 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SCHALL, Thomas, J.** [US/US]; 2050 Mill Ave., Menlo Park, CA 94025 (US). **TALBOT, Dale** [CA/US]; 67 Brentwood Avenue, San Francisco, CA 94127 (US). **MIAO, Zhenhua** [CN/US]; 7070 Rainbow Drive #2, San Jose, CA 95129 (US). **WEI, Zheng** [US/US]; 75 Duane Street #12, Redwood City, CA 94063 (US).

(74) Agents: **APPLE, Randolph, T. et al.**; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **TETHERED LIGANDS AND METHODS OF USE**




**Control (-)**



**ELC-Stalkokine (SK)**




**ELC-SK + soluble ELC**



**ELC-SK + soluble TECK**



**ELC-SK + soluble SLC**



**ELC-SK + soluble MCP-3**

(57) Abstract: The invention provides methods, reagents and devices for analysis of receptor ligand interactions, determining the profile of receptor expression in cells and cell populations, and diagnostic and drug screening methods. The invention makes use of immobilized tethered ligand fusion proteins having a ligand domain, a stalk domain, and optionally an immobilization domain.

WO 01/46698 A2

## TETHERED LIGANDS AND METHODS OF USE

5

### FIELD OF THE INVENTION

The invention relates to reagents and methods for characterizing receptor-ligand interactions, identifying modulators of such interactions, and characterizing the receptor expression profile of cells and cell populations. The invention finds application in the biomedical sciences.

10

### CROSS-REFERENCES TO RELATED APPLICATIONS

This application claims priority to U.S.S.N. 60/186626, filed March 3, 2000, the contents of which are incorporated by reference herein.

15

### BACKGROUND OF THE INVENTION

Communication between cells is a fundamental process involved in growth, differentiation, metabolism, and generation of biological responses (e.g., immune responses) in multicellular organisms. Often, cell-to-cell signaling is mediated by extracellular receptors or cell adhesion molecules. These membrane-associated molecules interact with other proteins, such as soluble factors such as peptide hormones, extracellular matrices, and cell surface molecules displayed by other cells. In some cases, signaling involves cell-to-cell contact (e.g., contact between two surface proteins on cells). An example of such an interaction is the binding of the T cell receptor on the surface of a T lymphocyte and the MHC/antigen complex on the surface of an antigen-presenting cell. A different type of cell-to-cell signaling is mediated by soluble polypeptides that, by interacting with a specific receptor on a target cell, lead to changes in target cell activity. An illustrative and important example cell-to-cell signaling mediated by soluble polypeptides is the activity of chemokines in mammalian systems.

30

Chemokines are a class of cytokines that play an important role in inflammatory responses, leukocyte trafficking, angiogenesis, and other biological processes related to the migration and activation of cells. As mediators of chemotaxis and inflammation, chemokines play roles in pathological conditions. For

example, the concentration of chemokine MCP-1 is higher in the synovial fluid of patients suffering from rheumatoid arthritis than that of patients suffering from other arthritic diseases.

Known chemokines are typically assigned to one of four subfamilies based on the arrangement of cysteine motifs. In the so-called alpha-chemokines, for example, the first two of four cysteines (starting from the amino terminus) are separated by an intervening amino acid (i.e., having the motif C-X-C). The beta-chemokines are characterized by the absence of an intervening amino acid between first two cysteines (i.e., comprising the motif C-C). The smaller gamma-chemokine family is characterized by a single C residue (gamma). The delta-chemokine family is characterized by a pair of cysteines separated by three residues (i.e., having the motif CX<sub>3</sub>C). The sole CX<sub>3</sub>C chemokine (fractalkine) is a type 1 membrane protein containing a chemokine domain tethered on a long mucin-like stalk. Fractalkine, also named neurotactin, has contains a chemokine domain at the amino terminus tethered on a long mucin-like stalk. For a recent review on chemokines, see Ward et al., 1998, *Immunity* 9:1-11 and Baggiolini et al., 1998, *Nature* 392:565-568, and the references cited therein.

Some chemokine activities (e.g., promigratory effects) are mediated by binding to an array of cell surface receptors on the surface of target leukocytes. These receptors are of the seven transmembrane spanning, G protein coupled receptor class (alternately referred to as 7TM or GPCR). Several seven-transmembrane-domain G protein-coupled receptors for C-C chemokines have been cloned: a C-C chemokine receptor-1 which recognizes MIP-1 $\alpha$ , RANTES, MCP-2, MCP-3, and MIP-5 (Neote et al., 1993, *Cell*, 72:415-415); CCR2 which is a receptor for MCP1, 2, 3 and 4 or 5; CCR3 which is a receptor for RANTES, MCP-2, 3, 4, MIP-5 and eotaxin; CCR5 which is a receptor for MIP-1 $\alpha$ , MIP-1 $\beta$  and RANTES; CCR4 which is a receptor for CMDC or TARC; CCR6 which is a receptor for LARC; and CCR7 which is a receptor for SLC and MIP-3 $\beta$  (reviewed in Sallusto et al., 1998, *Immunol. Today* 19:568 and Ward et al., 1998, *Immunity* 9:1-11).

Due to the importance of interactions between receptors (such as chemokine receptors) and their ligands in biological function, a need exists for rapid and effective methods for the characterization of such interactions.

### SUMMARY OF THE INVENTION

In one aspect the invention provides an assay device having a plurality of different immobilized tethered ligand fusion proteins organized in an array, where the ligand-stalk fusion proteins include a ligand domain and a stalk domain ("assay device"). In one embodiment, the ligand-stalk fusion proteins have a ligand domain, an intermediate stalk domain, and an immobilization domain ("assay device"). In an embodiment, the ligand domain and the stalk domain are not associated in a naturally occurring protein. In various embodiments, the immobilized tethered ligand fusion proteins include a mucin-derived stalk sequence, a fractalkine mucin repeat region sequence, and/or a ligand domain encoding a chemokine.

In another aspect, the invention provides a tethered ligand fusion protein, where the ligand domain is other than a chemokine sequence.

In another aspect, the invention provides a method for identifying an interaction between a receptor and a ligand by contacting a cell expressing a receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein and, detecting binding of the cell and the tethered ligand fusion protein, where binding of the cell to the fusion protein is correlated with an interaction between the receptor and the ligand corresponding to the ligand domain of the fusion protein. In an embodiment, the tethered ligand is immobilized on an assay device, supra. In various embodiments, the cell expresses a recombinant receptor (e.g., an orphan receptor). In an embodiment, the cell expresses a chemokine receptor. In one embodiment, an interaction between a receptor and more than one ligand is detected.

In another aspect, the invention provides a method for identifying a modulator of an interaction between a receptor and a ligand by contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the absence of a test compound, and measuring binding of the cells to an immobilized tethered ligand fusion protein, contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the presence of a test compound, and measuring binding of the cells to the immobilized tethered ligand fusion protein; and comparing the levels of binding, where decreased binding of cells in the presence of the test compound indicates that the test compound is an inhibitor of the interaction between the receptor and the ligand corresponding the ligand domain of said fusion protein,

and where increased binding of cells in the presence of the test compound indicates that the test compound is an enhancer of the interaction between the receptor and the receptor and the ligand corresponding the ligand domain of the fusion protein. In an embodiment, the tethered ligand is immobilized on an assay device, supra. In  
5 one embodiment, the receptor is a chemokine receptor.

In another aspect, the invention provides a method for detecting a profile of receptor expression in cells in a population of cells by contacting the population of cells with an immobilized tethered ligand fusion protein and detecting binding of cells of the population to the tethered ligand fusion protein, where binding  
10 of a cell in the cell population to a tethered ligand fusion protein is correlated with expression of a receptor that binds the ligand corresponding to the ligand domain of the fusion protein. In an embodiment, the tethered ligand is immobilized on an assay device, supra. In an embodiment, the contacting step involves contacting the population with a plurality of different fusion proteins and the detecting step involves  
15 detecting binding of cells to zero, one or more than one fusion proteins. In one embodiment, the population is heterogeneous, e.g., the population is obtained from synovial fluid, cerebral-spinal fluid, bronchial alveolar lavage (BAL) fluid, or blood. In an embodiment, the method includes quantitating the level of binding to each tethered ligand fusion protein in the array, or characterizing the cells bound at each  
20 sector of the array (e.g., by immunostaining).

In another aspect, the invention provides a method for diagnosis. The method includes the steps of obtaining a population of cells from a patient suspected of suffering from a disease, determining a receptor profile for the population, and, comparing said receptor profile with a profile characteristic of the disease state. In  
25 one embodiment the receptor profile is determined by contacting the population of cells with an immobilized tethered ligand fusion protein and identifying a tethered ligand fusion protein bound by cells of said population thereby identifying a receptor profile. In an embodiment, the tethered ligand is immobilized on an assay device, supra. In various embodiments, determining the method also includes quantitating  
30 the binding of cells at each sector of the array or characterizing the cells bound at each sector of the array, e.g., by immunostaining. In various embodiments, the disease is an inflammatory or allergic disease, or an autoimmune disease. In various embodiments, the population is obtained from synovial fluid, cerebral-spinal fluid, bronchial alveolar lavage (BAL) fluid, or blood.

In another aspect, the invention provides a method for detecting an effect of a drug or treatment on a patient by determining the receptor profile of a population of cells from the patient for the first time, administering the drug or treatment to the patient, determining the receptor profile of a population of cells from the patient for a second time, and comparing the receptor profiles obtained to determine the effect of the drug or treatment receptor-expressing cells in the patient. In an embodiment, the determining is carried out by contacting said population of cells with an immobilized tethered ligand fusion protein and identifying a subset of arrayed tethered ligand fusion proteins bound by cells of said population thereby identifying a receptor profile. In an embodiment, the tethered ligand is immobilized on an assay device, supra. In various embodiments, determining the method also includes quantitating the binding of cells at each sector of the array or characterizing the cells bound at each sector of the array, e.g., by immunostaining. In various embodiments, the population is obtained from synovial fluid, cerebral-spinal fluid, bronchial alveolar lavage (BAL) fluid, or blood.

In another aspect, the invention provides a method for identifying a modulator of an interaction between a receptor and a ligand by contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the presence of a test compound and determining the level of binding to the fusion protein, contacting cells expressing the receptor or a ligand-binding portion thereof with the immobilized tethered ligand fusion protein in the absence of a test compound; and determining the level of binding to the fusion protein, and comparing the level of binding, where the ligand-stalk fusion proteins include a ligand domain, an intermediate stalk domain, and, an immobilization domain and where decreased binding of cells in the presence of the test compound indicates that the test compound is an inhibitor of the interaction between the receptor and the ligand corresponding the ligand domain of fusion protein, and where increased binding of cells in the presence of the test compound indicates that the test compound is an enhancer of the interaction between the receptor and the ligand corresponding the ligand domain of the fusion protein. In one embodiment, the ligand domain and the stalk domain are not associated in a naturally occurring protein. In an embodiment, the stalk domain is carboxy-terminal to the ligand domain, and the immobilization domain is carboxy-terminal to the stalk domain.

In another aspect, the invention provides the use of tethered ligand fusion proteins in any of the methods described herein.

### BRIEF DESCRIPTION OF THE FIGURES

5           Figure 1 shows the structure of a tethered ligand.

Figure 2 shows the nucleotide sequence for a human CCR10 (SEQ ID NO:1) and the predicted amino acid sequence of the human CCR10 polypeptide (SEQ ID NO:2).

10

Figure 3 shows the identification of CCR10 ligands by adhesion to stalkokines. Figure 3A shows interrogation of immobilized stalkokine (SK) by HEK293-CCR10 cells, where 'control' shows background adhesion of HEK293-CCR10 cells to wells containing no stalkokine (anchoring antibodies alone are present); "ELC-stalkokine (SK)" shows strong adhesion of HEK293-CCR10 cells to locations containing ELC-stalkokines immobilized via anchoring antibodies; "ELC-SK + soluble ELC", "soluble TECK", or "soluble SLC" shows ablation of adhesion in the presence of excess concentrations of soluble recombinant 'native form' chemokines as shown; "ELC-SK + soluble MCP-3" shows no diminution in adhesion in the presence MCP-3 as representative of many non-competing chemokines. Wild type HEK293 cells showed no adhesion to any of the sites (not shown). Figure 3B shows the quantitation of adhesion of HEK293-CCR10 cells to ELC-stalkokine in the absence and presence of soluble chemokines from a representative experiment. Figure 3C shows the results of homologous competition binding assay using radiolabeled ELC in the presence of increasing concentrations of cold ELC on either HEK293-CCR10 cells (filled squares) or wild type HEK293 cells (open squares).

15  
20  
25

Figure 4 shows a schematic diagram of an expression plasmid encoding an ELC-stalkokine. Not all details of the plasmid are shown (e.g., pA site, polylinker, Ori, selection markers are not shown).

30

Figure 5 shows Western blot analysis of 32 different stalkokines. Each individual stalkokine construct was transfected into 293T and supernatants

containing the stalkokines were subjected to electrophoresis followed by Western blot analysis using the anti-poly histidine antibody. Equivalent volumes of supernatant were analyzed in each lane. Expression levels varied from a very robust accumulation in the case of MCP3 to a low level accumulation (IL-8).

Enhanced stalkokine accumulation can be achieved by addition of protease inhibitors. Due to glycosylation, the apparent molecular weight of the soluble stalkokines increased from the typical 35 kDa to around 90 kDa. Partial glycosylation resulted in an intermediate molecular weight, and it demonstrated by the very broad band or smear of signal. MW markers = 97.4, 68, 43, 29, 18.4, 14.3 kDa.

Figure 6 shows a displacement assay showing competition by a stalkokine with binding of a radiolabeled tracer chemokine to its cognate receptor.

Figure 7 shows a calcium mobilization assay using CEM cells expressing CCR4 receptors contacted with TARC-Stalkokine or MDC-Stalkokine, showing induction of calcium signaling in a receptor dependent manner.

## DETAILED DESCRIPTION

### 1. DEFINITIONS

The following definitions are provided to assist the reader in the practice of the invention. As used herein:

The term "receptor" has the ordinary meaning in the art and refers to a protein that specifically binds a second protein. Usually (i.e., in some embodiments) receptors are membrane associated proteins characterized by a transmembrane domain, an extracellular domain and an intracellular domain. Such receptors are called "membrane associated receptors." In other embodiments, the term "receptor" is used to refer to other types of polypeptide binding proteins, e.g., adhesion molecules, soluble binding proteins. It will be apparent that, as used herein, the term "receptor" does not refer to immunoglobulins.

The term "ligand," refers to peptides or polypeptide that bind to a receptor protein. Typically, ligands are involved in cell-to-cell signaling or adhesion of cells to other cells or extracellular matrix.



The term "specific binding" refers to binding between two molecules, for example, a ligand and a receptor, characterized by the ability of a protein (ligand) to associate with another specific molecule (receptor) even in the presence of many other diverse molecules, i.e., to show preferential binding of one molecule for another in a heterogeneous mixture of molecules. Specific binding of a ligand to a receptor is also evidenced by reduced binding of a detectably labeled ligand to the receptor in the presence of excess unlabeled ligand (i.e., a binding competition assay).

The term "orphan receptor" refers to a putative receptor polypeptide for which a naturally occurring cognate ligand is not known. Typically, orphan receptors are identified based on homology (sequence identity) to known receptors based on protein or gene sequence and structure.

The term "fusion protein," as used herein, refers to a composite polypeptide, i.e., a single contiguous amino acid sequence, made up of two or more (e.g., three) distinct polypeptides which are not normally fused together in a single amino acid sequence. Fusion proteins can generally be prepared using either recombinant nucleic acid methods, i.e., as a result of transcription and translation of a recombinant gene fusion product, which fusion comprises a segment encoding a first polypeptide region and a segment encoding a second polypeptide region, or by chemical synthesis methods well known in the art. Thus, for example, a single naturally occurring protein (e.g., Fractalkine) is not a fusion protein.

The term "recombinant" when used with reference to a cell indicates that the cell replicates a heterologous nucleic acid, or expresses a peptide or protein encoded by a heterologous nucleic acid. Recombinant cells can contain genes that are not found within the native (non-recombinant) form of the cell. Recombinant cells can also contain genes found in the native form of the cell wherein the genes are modified and re-introduced into the cell by artificial means (e.g., to increase expression of a receptor). The term also encompasses cells that contain a nucleic acid endogenous to the cell that has been modified without removing the nucleic acid from the cell; such modifications include those obtained by gene replacement, site-specific mutation, and related techniques.

A "recombinant expression cassette" or simply an "expression cassette" is a nucleic acid construct, generated recombinantly or synthetically, with nucleic acid elements that are capable of effecting expression of a structural gene in

hosts compatible with such sequences. Expression cassettes include at least promoters and optionally, transcription termination signals. Typically, the recombinant expression cassette includes a nucleic acid to be transcribed (e.g., a nucleic acid encoding a desired polypeptide), and a promoter. Additional factors  
5 necessary or helpful in effecting expression may also be used as described herein. Transcription termination signals, enhancers, and other nucleic acid sequences that influence gene expression, can also be included in an expression cassette. An "expression vector" is a vector that contains the necessary elements for the transcription and translation of the inserted coding sequence required for the  
10 expression system employed (e.g., a vector containing an expression cassette).

A "recombinant polynucleotide" or a "recombinant polypeptide" is a non-naturally occurring polynucleotide or polypeptide that includes nucleic acid or amino acid sequences, respectively, from more than one source nucleic acid or polypeptide.

15 The term "chemokine" (abbreviated "CK") refers to a class of cytokines involved in inflammatory responses, leukocyte trafficking, angiogenesis, and other biological processes related to the migration and activation of cells. Known chemokines are typically assigned to one of four subfamilies ( $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ ) based on the arrangement of cysteine motifs. A nonexhaustive list of chemokines is  
20 described in Table 2.

The term "receptor-bindable portion" when referring to a polypeptide ligand, refers to the portion of the ligand, often less than full-length naturally occurring protein) sufficient for binding to a cognate receptor (i.e., a receptor that binds the naturally occurring ligand).

## 25 2. INTRODUCTION

The present invention provides reagents, devices, and methods useful for analysis of protein-protein interactions such as receptor-ligand interactions. The invention provides reagents and methods for identification of modulators of such  
30 receptor-ligand interactions, and other methods, as described herein. In one embodiment, the invention is used to identify and characterize interactions between chemokines, e.g., human chemokines, and known or orphan chemokine receptors.

In an aspect of the invention, as described in detail *infra*, cells

expressing receptors are contacted with one or more tethered ligands and specific binding, if present, is detected and analyzed. Such contacting, detection and analysis is sometimes referred to as "interrogation." The "tethered ligands" used according to the invention are fusion proteins having at least two, and usually at least three, distinct domains: a ligand domain, a stalk domain, and, optionally, an immobilization domain. The cells may be recombinant cells expressing a predefined receptor protein (including cells expressing an orphan receptor), or may be cells isolated from a biological source (such as a tissue from a patient or non-human animal).

Assays in which binding of a cell expressing a receptor and the tethered ligand fusion proteins of the invention is detected may be carried out in a variety of ways.

In one embodiment, the interaction between one or more tethered ligands and one or more receptor expressing cells is detected and characterized. In a particular, the effects of one or more test compound(s) on the interaction can be assayed.

For example, tethered ligand fusion protein of the invention (i.e., a single species) may be immobilized on a surface (e.g., a solid surface such as a slide, plate, bead, dipstick, or the like) and contacted with a test cell or cell population.

In another embodiment of the invention, a plurality (i.e., at least 3, typically at least 10, often at least 15) of different tethered ligands (i.e., having different ligand domains) are immobilized in an array. The array can be interrogated by recombinant cells expressing receptors known or suspected of specifically binding the ligand domain of a tethered ligand. The effect of test compounds on the interaction of the cells and tethered ligands of the array can be determined. The array can also be interrogated by populations of naturally occurring cells, and the receptor profile of the population determined (for diagnostic, screening, or other purposes). Notably, the tethered arrays of the invention are useful for determining the ligand binding profile of orphan receptors.

The invention finds use in the characterization of a variety of protein-protein interactions. In particular, the invention is useful for the characterization of interactions between 7-transmembrane receptors (e.g., chemokine receptors) and putative ligands (e.g., chemokines). However, as will be apparent *infra*, the

invention is not limited to characterization of 7-transmembrane receptors.

### 3. TETHERED LIGANDS

As noted *supra*, the methods and devices of the invention make use of  
5 "tethered ligands." Tethered ligands are fusion proteins having a ligand domain, a  
stalk domain, and, usually, an immobilization domain. The stalk domain is  
positioned in between the ligand domain and the immobilization domain (i.e., it is  
"intermediate"). The stalk domain sequence may be contiguous with one or both of  
10 the other domains, but typically at least some additional (e.g., linker) sequences will  
be interposed. Typically, the ligand domain is located at the amino-terminus of the  
fusion protein, and the immobilization domain is at the carboxy-terminus (with the  
stalk domain in between). Fig. 1 shows a diagram of an exemplary tethered ligand.

In alternative embodiments, the ligand domain may be located at the  
15 carboxy-terminus of the fusion protein, and the immobilization domain at the amino-  
terminus. The latter orientation is useful when the ligand domain has the sequence  
of an intracellular protein, or intracellular domain of a transmembrane protein.

Techniques used in production of the tethered ligands of the invention  
(e.g., cloning and expression of recombinant proteins, including fusion proteins) are  
well known in the art and are described, for example, in Sambrook et al., 1989,  
20 Molecular Cloning: A Laboratory Manual, 2nd Ed., Vols. 1-3, Cold Spring Harbor  
Laboratory and Ausubel et al., Current Protocols In Molecular Biology, Greene  
Publishing and Wiley-Interscience, New York, including 1999 and 2000  
supplements. For example, it is understood that the various (e.g., at least 2) amino  
acid coding segments used to encode the fusion protein will be cloned in frame  
25 relative to each other, to encode the fusion protein.

The structure and production of tethered ligands will now be described  
in greater detail.

#### 3.1 Ligand Domain

30 The ligand domain of tethered ligands has the sequence of a protein or  
peptide ligand of interest, which may be an extracellular ligand or an intracellular  
ligand.

Extracellular ligands are those polypeptide ligands that are naturally  
bound by extracellular domains of membrane associated receptors; exemplary

extracellular ligands include chemokines (bound by chemokine receptors); other cytokines, interferon (bound by interferon receptors), dopamine (bound by dopamine receptors). Exemplary extracellular receptors are listed in Table 1, although many others will be apparent to those of skill. A particularly useful class of tethered ligands are those having a ligand domain corresponding to a chemokine. A tethered ligand having a chemokine domain is sometimes referred to as a "stalkokine," e.g., "TARC-stalkokine" for a tethered ligand having a TARC sequence in the ligand domain.

Chemokines and cytokines are well known in the art. Exemplary cytokines are listed in Table 2. References describing these and other cytokines are provided in the R&D Systems Catalog (1999) and (2000) R&D Systems Inc., 614 McKinley Place N.E. MN 55413, the R&D online catalog at <http://www.rndsystems.com> (e.g., October 10, 1999), both of which are incorporated by reference for all purposes; the CFB (Cytokine Facts Book, 1994, Academic Press Ltd.), Chemokine Facts Book, 1997, Academic Press Ltd., incorporated by reference for all purposes, and the GenBank protein sequence database (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi>).

It will be appreciated that a ligand domain need not necessarily include the complete sequence of the intracellular or extracellular ligand to which it corresponds; in various embodiments, the ligand domain will comprise at least a receptor-bindable portion of the naturally occurring ligand.

**Table 1**

**Exemplary Receptors**

Receptor Class	Subclasses & Examples	GenBank Accession Number
Chemokine Receptors	1. CCR1 2. CCR2 3. CCR3 4. CCR4 5. CCR5 6. CCR6 7. CCR7 8. CCR8 9. CCR9 10. CCR10 11. CCR11 12. CXCR1 (IL-8R alpha) 13. CXCR2 (IL-8R beta) 14. CXCR3 (GPR9)	NM001295 NM000647 NM001837 AB023892 NM000579 NM004367 NM001838 NM005201 NM006641 AF233281 NM016602 L19591 NM001557 NM001504

	15. CXCR4 16. CXCR5 17. XCR1 18. CX <sub>3</sub> CR1 19. CMV US28 20. HHV8ORF74 21. Duffy Ag	NM003467 AJ002211 NM005283 U20350 L20501 AF179931 AF030521
Other Receptor Classes	Peptide Angiotensin Angiotensin type 1 Angiotensin type 2 Bombesin Bradykinin C5a anaphylatoxin Fmet-leu-phe Interleukin-8 Interleukin-8 type A Interleukin-8 type B Endothelin Melanocortin Melanocyte stimulating hormone Adrenocorticotrophic hormone Melanocortin hormone Neuropeptide Y Neurotensin Opioid Opioid type D Opioid type K Opioid type M Opioid type X Somatostatin Somatostatin type 1 Somatostatin type 2 Somatostatin type 3 Somatostatin type 4 Somatostatin type 5 Tachykinin Substance P (NK1) Substance K (NK2) Neuromedin K (NK3) Thrombin Vasopressin-like Vasopressin Oxytocin Galanin Proteinase activated Orexin Chemokine/chemotactic factors like Hormone protein Follicle stimulating hormone Lutropin-choriogonadotropic hormone Thyrotropin Viral Class A Orphan/other Burkitt's lymphoma receptor Mas proto-oncogene RDC1 EDG GPR	

	RBS11 BONZO Chemokine receptor-like 2 G10D GP40 like APJ like
	Growth Factor Receptors Hormone Receptors Class I cytokine receptors (hematopoietin family receptors)
	IL-6 (sharing gp 130) IL-6 IL-11 CNTF LIF Oncostatin M Cardiotrophin 1
	GM-CSF (sharing $\beta$ chain) GM-CSF IL-3 IL-5
	IL-2 (sharing $\gamma$ chain) IL-2 IL-4 IL-7 IL-9 IL-15
	IL-13 (sharing $\alpha$ chain) IL-13 IL-14
	IL-12 G-CSF Erythropoietin Growth hormone Prolactin
	IFN- $\alpha/\beta$ family IFN- $\gamma$ IL-10
	TNF- $\alpha$ TNF- $\beta$ (LT- $\alpha$ ) LT- $\alpha$ / LT- $\beta$ heteromes NGF Fas Ligand CD40 ligand TRAIL
	IL-1 $\alpha$ IL-1 $\beta$ IL-1 receptor antagonist
	TGF- $\beta$ Bone morphogenetic proteins

Table 2  
Exemplary Cytokines

1. TNF-alpha	39. Activins	78. IL-13
2. CD27L	40. TRAIL	79. TWEAK
3. TGF-beta	41. TALL-1	80. GITRL
4. TNF-beta	42. IL-15	81. Lymphotoxin
5. CD30L	43. IL-5	82. BRAK
6. IL-2	44. IL-1beta	83. PARC
7. VEGF (VEGF-A)	45. IL-10	84. Eotaxin
8. EGF	46. IGF-II	85. Eotaxin-2
9. HB-EGF	47. MSP	86. MPIF-1
10. IL-6	48. FGF-basic	87. HCC-1
11. SCF	49. BDNF	88. HCC-4
12. BMPs	50. Erythropoietin	89. SCYA26
13. Lymphotoxin-beta	51. PDGF-AB	90. MIP-1alpha
14. CD40L	52. VEGF-D	91. MIP-1beta
15. IL-7	53. Betacellulin	92. MIP-1delta
16. VEGF-B	54. G-CSF	93. I-309
17. TGF-alpha	55. CNTF	94. 6-Ckine
18. SMDF	56. Midkine	95. RANTES
19. LIF	57. Inhibins	96. MCP-1
20. Flt-3 Ligand	58. TRANCE	97. MCP-2
21. GDNFs	59. 4-1BBL	98. MCP-3
22. Fas Ligand	60. IL-4	99. MCP-4
23. April	61. GM-CSF	100. TARC
24. IL-9	62. IL-18	101. MIP-3alpha
25. IL-3	63. IL-17	102. MIP-3beta
26. IL-1alpha	64. IL-16	103. TECK
27. IFN-gamma	65. IL-12	104. MDC
28. IGF-I	66. FGF-3-19	105. IL-8
29. HGF	67. Neurotrophins	106. GCP-2
30. FGF-acidic	68. Angiopoietins (1-4)	107. MIG
31. NGF	69. PDGF-BB	108. SDF-1
32. Thrombopoietin	70. PIGF	109. I-TAC
33. PDGF-AA	71. Heregulins	110. PF4
34. VEGF-C	72. Leptin	111. ENA-78
35. Amphiregulin	73. Oncostatin M	112. IP-10
36. IL-11	74. Pleiotrophin	113. NAP-2
37. Cardiotrophin-1	75. MIS	114. GRO
38. M-CSF	76. LIGHT	115. BLC/BCA-1
	77. OX40L	

In some alternative embodiments, the "receptor" is not the extracellular domain of a membrane associated receptor, but a different protein binding moiety, such as the protein-binding domain of an intracellular protein or protein domain. For



example, the "receptor-ligand pair" used in the invention can be the intracellular domain of a CCR1 ("receptor") and a G-protein ("ligand"). In this case, the "ligand" is the G-protein domain and is expressed as a fusion protein with a stalk domain (e.g., a fractalkine mucin region sequence). The "receptor" is the intracellular domain of CCR1, that can be expressed as an amino-terminal fusion protein with a single transmembrane receptor (such as EGF receptor with a deleted kinase domain). This allows the CCR1 intracellular domain, for example, to be expressed extracellularly (i.e. on the surface of a cell). Thus, by immobilizing the G-protein tethered ligand onto a solid surface one can interrogate the cell expressing the CCR1 intracellular domain. Another example of an intracellular ligand (a protein that binds an intracellular protein, or intracellular protein domain), is a G protein that binds a G-protein coupled glutamate receptor.

### 3.2 Stalk Domain

The stalk domain of the tethered ligand fusion protein functions, in part, to display the ligand domain by elevating it a considerable distance above the substrate to which the tethered ligand fusion protein is immobilized. Without intending to be bound by a particular mechanism, it is believed that displaying the ligand a significant distance above the surface increases binding the ligand with the receptor displayed in receptor-expressing cells.

Typically, the length of the stalk domain of the fusion protein is at least about 50 amino acid residues at least about 75 amino acid residues, at least about 100 amino acid residues, often at least about 150 residues, and frequently at least about 200 residues. Typically, the stalk domain is between about 200 residues and 500 residues, and usually between about 200 residues and 300 residues. Generally, the ligand domain of the polypeptide is displayed at least about 20nm (e.g., between about 20 nm and about 60 nm), or at least about 30 nm away from (i.e., above) the surface to which the tethered ligand is immobilized. In embodiments, the ligand domain peptide is displayed at between around 30-40 nm above the surface. The measurement of the extension of the ligand domain is typically done by electron microscopy after heavy metal shadowing (see, e.g., Fong et al, 2000, *J Exp Med* 188:1413-19).

In addition to elevating the ligand domain away from the surface, the flexibility of the stalk region is believed to permit the ligand domain to adopt a variety

of orientations, increasing the likelihood of a strong interaction with the receptor, and having unexpected advantages compared to other methods of immobilizing or displaying a ligand, such as standard ELISA. Typically, the stalk domain is selected to have sufficient rigidity to elevate the ligand but has flexibility to allow the ligand domain to adopt a variety of orientations.

In one embodiment, the stalk domain is derived from the fractalkine polypeptide (Bazan et al., 1997, *Nature* 385:640-644; See also, WO 97/27299). Fractalkine is a naturally occurring type 1 membrane protein containing a chemokine domain tethered on a long mucin-like stalk. Human fractalkine cDNA (Genbank Accession No. U84487) encodes a 397 residue membrane protein with a 24 residue predicted signal peptide, a 76 residue chemokine domain, a 241 residue stalk region containing 17 degenerate mucin-like repeats rich in motifs for O-glycosylated serine and threonine residues, a 19 residue transmembrane segment and a 37 residue cytoplasmic domain.

Thus, in exemplary embodiments, the stalk region has the sequence provided in Table 3A or 3B. In other embodiments, the stalk region includes at least one mucin repeat segment from a fractalkine stalk region, such as the human sequence shown in Table 4. In related embodiments, the stalk region has at least 10, at least 25, or at least 50 contiguous residues of a fractalkine stalk region, such as that shown in Table 3A or 3B, (e.g., subsequences or interactions thereof).

Table 3A

Human Fractalkine Mucin-Repeat Region

IGEVKPRRTTPAAGGMDES VVLEPEATGESSSLEPTPSSQEAQRALGTSP ELPTGVT  
GSSGTRL PPTPKAQDGGPVGTELFRVPPVSTAATWQSSAPHQPGPSLWAEAKTS  
EAPSTQDPSTQASTASSPAPEENAPSEGQRVWGQGQSPRPENSLEREEMGPVP  
AHTDAFQDWGPGSMAHVSVPVSSEGTPSREPVASGSWTPKAE EPIHATMDPQR  
LGVLT PVP (SEQ ID NO:3)

Table 3B

Human Fractalkine Mucin-Repeat Sequence Residues 100-336

GGTFEKQIGEVKPRRTTPAAGGMDES VVLEPEATGESSSLEPTPSSQEAQRALGTS  
PELPTGVTGSSGTRL PPTPKAQDGGPVGTELFRVPPVSTAATWQSSAPHQPGPSL  
WAEAKTSEAPSTQDPSTQASTASSPAPEENAPSEGQRVWGQGQSPRPENSLERE  
EMGPVPAHTDAFQDWGPGSMAHVSVPVSSEGTPSREPVASGSWTPKAE EPIHA  
TMDPQRLGVLT PVPDAQA (SEQ ID NO:4)

Table 4  
Mucin Repeat Domains

	1. IGEVKPRTTP	(SEQ ID NO:5)
	2. GGMDSEVVLEP	(SEQ ID NO:6)
5	3. TGESSSLEPTP	(SEQ ID NO:7)
	4. LGTSPELPTG	(SEQ ID NO:8)
	5. TGSSGTRLPPPTP	(SEQ ID NO:9)
	6. VGTELFVRVPPVS	(SEQ ID NO:10)
	7. AATWQSSAPHQ	(SEQ ID NO:11)
10	8. PGPSLWAEAKTS	(SEQ ID NO:12)
	9. EAPSTQDPST	(SEQ ID NO:13)
	10. QASTASSPAP	(SEQ ID NO:14)
	11. VWGQGQSPRP	(SEQ ID NO:15)
	12. SLEREEMGPVP	(SEQ ID NO:16)
15	13. AHTDAFQDWG	(SEQ ID NO:17)
	14. PGSMAHVSVVP	(SEQ ID NO:18)
	15. EGTPSREPVA	(SEQ ID NO:19)
	16. SGSWTPKAEPP	(SEQ ID NO:20)
20	17. QRLGVLITPVP	(SEQ ID NO:21)

In related embodiments, the stalk region of the tethered ligand has a sequence from a fractalkine of a non-human species such as mouse (see, Lloyd et al, 1997, *Nature* 387:611-617; Genbank Accession No. AF010586) and other mammals (e.g., porcine, bovine, ovine, rat, rabbit, and non-human primate mammals).

In a related embodiment, the stalk domain is derived from other mucin family members, such as MUC-type mucins. MUC-type mucins are a family of structurally related molecules that are heavily glycosylated and are expressed in epithelia of the respiratory, gastrointestinal, and reproductive tracts, e.g., MUC1 (GenBank accession number AF125525), MUC2 (L21998), MUC3 (AF113616), MUC4 (AJ000281), MUC5AC (U83139), MUC5B (AJ001402), MUC6 (U97698), MUC7 (L13283), MUC8 (U14383), MUC9 (oviductin) (AW271430). In other embodiments, the stalk domain has sequences from MAdCAM-1, GlyCAM-1, CD34 (See, e.g., Girard & Springer 1995; Van Klinken et al., 1988, *Anal Biochem* 265:103-16), consensus repeats from E- selectin, P- selectin, or L-selectin, or viral glycoprotein spikes (e.g., glycoproteins of viral origin, such as influenza, herpes simplex, human immunodeficiency, or Tobacco mosaic virus) In particular, the influenza virus neuraminidase protein (accession number 091744), and especially the hypervariable stalk region from amino acids 36 through 90 inclusive (i.e., comprising

HFKQYECSSPPNNQVIPCQPTIERNITEIVYLTNTTIEKEICPKLVEYRNWSKP (SEQ ID NO:22) and concatenations thereof), are useful for a stalk to display immobilized ligands:

Additional stalk sequences can be tested (for suitability for use in the methods of the invention) using the assay described in Example 1, *infra*, but substituting the new stalk sequence for the fractalkine domain of Example 1. Briefly, to identify new suitable stalk region sequence, sequences encoding potential display stalks are cloned into an expression vector [e.g., pcDNA3.1/Myc-His(-)B (Invitrogen<sup>TM</sup> Corp, Carlsbad CA) or other suitable vector (e.g., pcDNA3.1/Myc-His(-)A/C (Invitrogen<sup>TM</sup> Corp.) or a similar vector], for example as an *EcoRI*-linked PCR fragment to form a fusion protein having an ELC ligand domain at the amino terminus and a 6xHis sequence at the carboxy-terminus. That is, the stalk-encoding sequence is inserted upstream in an analogous position as was used for fractalkine domain of the ELC-stalkokine (i.e., between an ELC binding motif and a poly-histidine immobilization domain. The resulting plasmid is expressed in mammalian cells (e.g., 293 cells) and the tethered ligand protein expressed, e.g., as described in Example 1. The tethered ligand is then assayed for ability to mediate binding of cells expressing CCR10, as described *infra*. Variations of this assay will be apparent to those of skill in the art.

### 3.3 Immobilization Domain

In various embodiments, the fusion protein has an immobilization domain to facilitate immobilization of the protein to a solid substrate. Often, immobilization domain is a short (i.e., fewer than 10 residues) epitope tag (i.e., a sequence recognized by an antibody, typically a monoclonal antibody) such as polyhistidine (Bush et al, 1991, *J. Biol Chem* 266:13811-14), SEAP (Berger et al, 1988, *Gene* 66:1-10), or M1 and M2 flag (see, e.g, U.S. Pat. Nos. 5,011,912; 4,851,341; 4,703,004; 4,782,137). In some embodiments of the invention, the tethered ligand does not have a separate immobilization domain. Instead, the stalk domain is directly conjugated to the substrate, the stalk domain is immobilized via an anti-stalk (i.e., mucin) sequence antibody, or via some other immobilization method.

### 3.4 Production of Tethered Ligands

As noted *supra*, methods for constructing and expressing fusion proteins are well known. Fusion proteins generally are described in Ausubel et al., *supra*, Kroll et al., 1993, *DNA Cell. Biol.* 12:441, and Imai et al., 1997, *Cell* 91:521-30.

The tethered ligand fusion proteins of the invention are typically made by (1) constructing a vector (e.g., plasmid, phage or phagemid) comprising a polynucleotide sequence encoding the desired polypeptide, (2) introducing the vector into a suitable expression system (e.g., a prokaryotic, insect, mammalian, or cell free expression system), (3) expressing the fusion protein and (4) optionally purifying the fusion protein.

(1) In one embodiment, expression of the tethered ligand fusion protein comprises inserting the coding sequence into an appropriate expression vector (i.e., a vector that contains the necessary elements for the transcription and translation of the inserted coding sequence required for the expression system employed, e.g., control elements including enhancers, promoters, transcription terminators, origins of replication, a suitable initiation codon (e.g., methionine), open reading frame, and translational regulatory signals (e.g., a ribosome binding site, a termination codon and a polyadenylation sequence. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used.

The coding sequence of the tethered ligand fusion protein includes ligand, stalk and immobilization domains as described elsewhere herein. Polynucleotides encoding the amino acid sequence for each domain can be obtained in a variety of ways known in the art; typically the polynucleotides are obtained by PCR amplification of cloned plasmids, cDNA libraries, and cDNA generated by reverse transcription of RNA, using primers designed based on sequences determined by the practitioner or, more often, publicly available (e.g., through GenBank). Typically, the primers include linker regions (e.g., sequences including restriction sites) to facilitate cloning and manipulation in production of the fusion construct. The polynucleotides corresponding to the ligand, stalk and immobilization regions are joined in-frame to produce the fusion protein-encoding sequence.

The tethered ligand proteins of the invention may be expressed as

secreted proteins or as nonsecreted proteins. Preferably, the fusion proteins are secreted. When the naturally occurring ligand includes a signal peptide (e.g., chemokine ligands) secretion is easily achieved by including the signal sequence encoding DNA in the fusion gene. Alternatively, and in the case of ligands not naturally secreted, a heterologous or artificial signal peptide is included in the fusion protein (see, e.g., Lui et al, 1993, *PNAS USA*, 90:8957-61).

(2) The tethered ligand fusion protein vectors may be introduced into a cell (e.g., bacterial, yeast, insect, and mammalian cells) by a variety of methods. The nucleic acid expression vectors (typically dsDNA) of the invention can be transferred into the chosen host cell by well-known methods such as calcium chloride transformation (for bacterial systems), electroporation, calcium phosphate treatment, liposome-mediated transformation, injection and microinjection, ballistic methods, virosomes, immunoliposomes, polycation:nucleic acid conjugates, naked DNA, artificial virions, fusion to the herpes virus structural protein VP22 (Elliot and O'Hare, *Cell* 88:223), agent-enhanced uptake of DNA, and other methods known in the art. See Ausubel, *supra*.

(3) A variety of expression systems suitable for expression of tethered ligands are known in the art. Useful bacterial expression systems include *E. coli*, bacilli (such as *Bacillus subtilis*), other enterobacteriaceae (such as *Salmonella*, *Serratia*, and various *Pseudomonas* species) or other bacterial hosts (e.g., *Streptococcus cremoris*, *Streptococcus lactis*, *Streptococcus thermophilus*, *Leuconostoc citrovorum*, *Leuconostoc mesenteroides*, *Lactobacillus acidophilus*, *Lactobacillus lactis*, *Bifidobacterium bifidum*, *Bifidobacterium breve*, and *Bifidobacterium longum*). The tethered ligand fusion protein expression constructs useful in prokaryotes include recombinant bacteriophage, plasmid or cosmid DNA expression vectors, or the like, and typically include promoter sequences. Illustrative promoters include inducible promoters, such as the lac promoter, the hybrid lacZ promoter of the Bluescript7 phagemid (Stratagene, La Jolla CA) or pSport1 (Gibco BRL); phage lambda promoter systems; a tryptophan (*trp*) promoter system; and *ptrp-lac* hybrids and the like. Bacterial expression constructs optionally include a ribosome binding site and transcription termination signal regulatory sequences. Illustrative examples of specific vectors useful for expression include, for example, pTrcHis2, (Invitrogen, San Diego CA),

When tethered ligand fusion protein is expressed in yeast, a number of

suitable vectors are available, including plasmid and yeast artificial chromosomes (YACs) vectors. The vectors typically include expression control sequences, such as constitutive or inducible promoters (e.g., such as alpha factor, alcohol oxidase, PGH, and 3-phosphoglycerate kinase or other glycolytic enzymes), and an origin of replication, termination sequences and the like, as desired. Suitable vectors for use in *Pichia* include pPICZ, His6/pPICZB, pPICZalpha, pPIC3.5K, pPIC9K, pA0815, pGAP2A, B & C, pGAP2alpha A, B, and C (Invitrogen, San Diego, CA) and numerous others known in the art or to be developed. In one embodiment, the vector His6/pPICZB (Invitrogen, San Diego, CA) is used to express a His6-tethered ligand fusion protein fusion protein in the yeast *Pichia pastoris*. An example of a vector useful in *Saccharomyces* is pYES2 (Invitrogen, San Diego, CA).

Another expression system provided by the invention for expression of tethered ligand fusion protein is an insect system. A preferred system uses a baculovirus polyhedrin promoter. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequence encoding the gene of interest may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the sequence, e.g., encoding the tethered ligand fusion protein, will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae, in which the tethered ligand fusion protein sequence is then expressed (see, for general methods, Smith et al., 1983, *J. Virol.*, 46:584; Engelhard et al., 1994, *Proc. Natl. Acad. Sci.* 91:3224-7). Useful vectors for baculovirus expression include pBlueBacHis2 A, B & C, pBlueBac4.5, pMelBacB and numerous others known in the art or to be developed. Illustrative examples of tethered ligand fusion protein expression constructs useful in insect cells are provided in Example 6, *infra*.

The present invention also provides expression systems in mammals and mammalian cells. As noted *supra*, tethered ligand fusion protein polynucleotides may be expressed in mammalian cells (e.g., human cells) for production of significant quantities of tethered ligand fusion protein polypeptides are well known in the art and include those described *infra* §5.1.1

Optimum expression times are conveniently determined by analysis of the secreted protein by Western blot analysis (e.g., using a common epitope tag, e.g., poly histidine) for detection.

(4) It is sometimes desirable to purify a secreted tethered ligand fusion protein from cell media, and generally necessary to purify nonsecreted proteins prior to use. Purification can be carried out using a variety of methods, including, for example, metal-chelate affinity chromatography of fusion proteins containing polyhistidine tracts (e.g. His<sub>6</sub>), protein A domains or fragments (which allow purification on immobilized immunoglobulin), and the domain utilized in the  
5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
1023



#### 4. IMMOBILIZATION AND PREPARATION OF ARRAYS

##### 4.1 Immobilization on Substrates

In some embodiments, the tethered ligands (e.g., stalkokines) of the invention are immobilized on a solid surface. The substrate to which tethered ligand is bound may be in any of a variety of forms, e.g., a microtiter dish, a test tube, a dipstick, a microcentrifuge tube, a bead, a spinnable disk, and the like. Suitable materials include glass, plastic (e.g., polyethylene, PVC, polypropylene, polystyrene, and the like), protein, paper, carbohydrate, and other solid supports. Other materials that may be employed include ceramics, metals, metalloids, semiconductive materials, cements and the like.

As noted *supra*, in some embodiments, the fusion proteins are organized as an array. The term "array," as used herein, refers to an ordered arrangement of immobilized fusion proteins, in which particular different fusion proteins (i.e., having different ligand domains) are located at different predetermined sites on the substrate. Because the location of particular fusion proteins on the array is known, binding of receptor-bearing cells to that location can be correlated with specific binding of the receptor or receptors displayed by the cell and the ligand (e.g., chemokine) domain of the fusion protein.

Immobilization of fusion proteins on beads (individually or in groups) is another particularly useful approach. In one embodiment, individual tethered ligand fusion proteins are immobilized on beads, allowed to bind to cells expressing cognate receptors, and the resulting complex is separated from unbound cells. In one embodiment, mixtures of distinguishable beads are used. Distinguishable beads are beads that can be separated from each other on the basis of a property such as size, magnetic property, color (e.g., using FACS) or affinity tag (e.g., a bead coated with protein A can be separated from a bead not coated with protein A by using IgG affinity methods). According to one embodiment, each distinguishable bead is associated with a species or particular combination of ligand fusion proteins. Cell binding to particular tethered ligand fusion proteins may be determined; similarly, the effect of test compounds (i.e., agonists and antagonists of binding) may be determined.

##### 4.2 Substrates and Methods for Immobilization

Methods for immobilizing proteins are well known in the art, and

include covalent and non-covalent methods. It will be appreciated that the choice of method will depend in part on the substrate and detection system selected.

#### 4.2.1 Non-covalent immobilization

5           One suitable immobilization method is antibody-mediated immobilization. According to this method, an antibody specific for the sequence of an "immobilization domain" of the tethered ligand is itself immobilized on the substrate (e.g., by adsorption). One advantage of this approach is that a single antibody may be adhered to the substrate and used for immobilization of a number  
10 of different tethered ligands (sharing the same immobilization domain). For example, an immobilization domain consisting of poly-histidine (Bush et al, 1991, *J. Biol Chem* 266:13811-14) can be bound by an anti-histidine monoclonal antibody (R&D Systems, Minneapolis, MN); an immobilization domain consisting of secreted alkaline phosphatase ("SEAP") (Berger et al, 1988, *Gene* 66:1-10) can be bound by  
15 anti-SEAP (Sigma Chemical Company, St. Louis, MO); an immobilization domain consisting of a FLAG epitope can be bound by anti-FLAG. Other ligand-antiligand immobilization methods are also suitable (e.g., an immobilization domain consisting of protein A sequences (Harlow and Lane, 1988, *ANTIBODIES A LABORATORY MANUAL*, Cold Spring Harbor Laboratory; Sigma Chemical Co., St. Louis, MO) can be bound  
20 by IgG; and an immobilization domain consisting of streptavidin can be bound by biotin (Harlow & Lane, *supra*; Sigma Chemical Co., St. Louis, MO).

When antibody-mediated immobilization methods are used, glass is an especially useful substrate (e.g., microscope quality glass slides). When arrays are desired, the glass substrates may be printed with a hydrophobic (e.g., Teflon) mask  
25 to form wells. Preprinted glass slides with 3, 10 and 21 wells per 14.5 cm<sup>2</sup> slide "working area" are available from, e.g., SPI Supplies, West Chester, PA; also see U.S.Pat. No. 4,011,350). In certain applications, a large format (12.4 cm x 8.3 cm) glass slide printed in a 96 well format is used; this format facilitates the use of automated liquid handling equipment and utilization of 96 well format plate readers of  
30 various types (fluorescent, colorimetric, scintillation). However, higher densities may be used (e.g., more than 1 tethered ligand per cm<sup>2</sup>). Often the tethered ligand array will include at least about 3, at least about 5, at least about 10, at least about 15, at least about 20 or more different ligands.

Typically, antibodies are bound to substrates (e.g., glass substrates) by adsorption. Suitable adsorption conditions are well known in the art and include incubation of 0.5-50 ug/ml (e.g., 10 ug/ml) mAb in buffer (e.g., PBS, or 50 to 300 mM Tris, MOPS, HEPES, PIPES, acetate buffers, pHs 6.5 to 8, at 4°C to 37°C and from 1hr to more than 24 hours.

Antibody-mediated immobilization may provide certain advantages over other methods for immobilization, due to the flexibility of the antibody hinge region, allowing the ligand domain to adopt additional orientations.

Usually, the fusion protein is immobilized prior to contacting with receptor-expressing cells, however, in some embodiments, the immobilization follows the contacting step.

#### 4.2.2 Covalent immobilization

As noted *supra*, the tethered ligand may be covalently bound or noncovalently attached through nonspecific bonding. If covalent bonding between a the fusion protein and the surface is desired, the surface will usually be polyfunctional or be capable of being polyfunctionalized. Functional groups which may be present on the surface and used for linking can include carboxylic acids, aldehydes, amino groups, cyano groups, ethylenic groups, hydroxyl groups, mercapto groups and the like. The manner of linking a wide variety of compounds to various surfaces is well known and is amply illustrated in the literature.

### 5. INTERROGATION OF STALKOKINES AND STALKOKINE ARRAYS

According to the methods of the invention, cells expressing receptors (e.g., chemokine receptors) are contacted with one or more tethered ligand fusion proteins (e.g., individual ligands or an array of different tethered ligands) under conditions in which binding between the ligand and cognate receptor (if present) occurs.

Typically, the binding conditions are physiological conditions under which receptor-ligand interactions occur or are expected to occur. Thus, in one embodiment, the cells are contacted with the tethered ligand in a buffered solution (e.g., PBS, TBS, etc).

Binding is typically allowed to proceed for between 0.5 and 10 h (typically between 0.5 and 3 h, e.g., 1.5 h) at 14°C to 37°C (e.g., room temperature),

with or without gentle agitation. Preferred cell concentrations for binding range from  $0.5\text{-}5 \times 10^6/\text{ml}$  (e.g.,  $1\text{-}2 \times 10^6/\text{ml}$ ), but concentrations outside this range may be used (e.g., in the case of rare cells from a patient sample).

5 In some embodiments, the cells are pretreated prior to incubation with the array or individual tethered ligands. For example, cells may be decorated with fluorescently labeled antibodies (e.g., anti-CD3 cell marker) or labeled intracellularly with fluorescent dyes such as Calcein (Molecular Probes, Eugene, OR), or labeled using radioisotopes. Such labeling facilitates quantitation of cell binding to tethered ligands (e.g., sites on an array) as well as characterization of the binding cells. For  
10 example, in an experiment analyzing chemokine receptors expressed on T lymphocytes, PBMCs may be isolated and T cells stained for with fluorescein labeled anti-CD4 and rhodamine labeled anti-CD8. Typically the cells are stained for 30 minutes, washed 3x in PBS, and resuspended at  $1 \times 10^6$  cells per ml in PBS. The cells are then contacted with a tethered ligand array described herein, and allowed  
15 to bind at room temperature for 1 hour. The slide is then washed to remove unbound cells, and the adherent cells detected using a fluorescent plate reader. The fluorescein tag and the rhodamine tag are separately detected, with co-localization identifying T cells.

By analysis of the tethered chemokine to which each cell type is  
20 bound, and the ratio of the two signals, the cell type being localized at a particular tethered ligand may be determined (i.e., the cell may be characterized and the cells quantitated). It will be apparent that by staining a heterogeneous population of cells (e.g., from a patient with multiple sclerosis or rheumatoid arthritis) subpopulations of cells expressing particular receptors, or combinations of receptors, can be identified,  
25 e.g., for diagnostic purposes. Similarly, this protocol is useful in drug development (e.g., screening for compounds that alter the binding pattern of individual T cells to the tethered chemokines).

### 5.1 Interrogation with recombinant cells expressing receptors

30 In one embodiment, the methods of the invention are practiced using recombinant cells expressing receptors (i.e., cells transiently or stably transfected with expression vectors encoding the receptor protein of interest).

### 5.1.1 Cells Expression Recombinant Receptor Proteins

The recombinantly expressed receptor proteins used in the assay may be any of a variety of binding proteins, such as seven transmembrane spanning, G protein coupled receptor class (e.g., chemokine receptors), receptor tyrosine kinases (e.g., EGF receptor, insulin receptor, IGF-1 receptor, NGF receptor, PDGF receptor, M-CSF receptor, FGF receptor, VEGF receptor) and receptors for any ligand or ligand class listed in Table 2, *supra*. Exemplary receptors are shown in Table 1.

Additional description of 7-transmembrane receptors, including links to sequence databases, is found at <http://swift.embl-heidelberg.de/7tm/phylo/phylo.html>.

Numerous additional protein binding receptors are described in the literature and/or in GenBank. Using this information, one of ordinary skill can prepare cells expressing a variety of receptors using routine techniques.

Transformed cell lines expressing receptors of interest and/or expression vectors encoding various receptors are known (see, e.g., American Type Culture Collection, 10801 University Boulevard Manassas, VA 20110-2209; <http://www.atcc.org/>). Alternatively, provided with the DNA sequence of the gene or cDNA encoding a receptor, the receptor may be expressed using routine methods for expression of recombinant proteins. It will be understood that it usually is not necessary to express a full-length or naturally occurring receptor sequence; a portion, variant or fragment capable of binding the ligand is sufficient.

Usually, the receptor-encoding sequence of interest is cloned into a "recombinant expression cassette" expression vector and introduced (e.g., transfected) into an appropriate host cell.

Although receptors may be expressed in a variety of cell types capable of expressing a membrane-bound receptor protein (e.g., insect, and the like), the receptors used in the present invention typically are expressed in mammalian cells.

Host cells useful for receptor expression include, but are not limited to, insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus), mammalian cells expression systems and the like. See, e.g., Ausubel et al., *supra*. In mammalian host cells, a number of expression systems may be utilized, e.g., constitutive expression vectors such as pCMV-myc (Clontech, Palo Alto CA), inducible expression system such as pTRE2 (Clontech, Palo Alto CA) (using tetracycline induction), vectors such as the bicistronic vector pIRES (Clontech, Palo Alto CA) capable of driving the receptor and a drug resistance

gene, and the like. In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific fashion desired, e.g., modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such host cells include, but are not limited to, monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293; Graham et al., *J. Gen. Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); CHO (ATCC CCL 61 and CRL 9618); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL 1587); human cervical carcinoma cells (HeLa, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather, et al., *Annals N.Y. Acad. Sci.* 383:44-46 (1982); MDCK cells (ATCC CCL 34 and CRL 6253); HEK 293 cells (ATCC CRL 1573); and WI-38 cells (ATCC CCL 75; ATCC: American Type Culture Collection, Rockville, MD). The use of mammalian tissue cell culture to express polypeptides is discussed generally in Winnacker, *FROM GENES TO CLONES* (VCH Publishers, N.Y., N.Y., 1987).

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express receptors may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with the receptor encoding DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. When appropriate, codon usage may be modified for expression in non-human cells or non-mammalian cells. Following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched medium, and then switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which

express the receptor proteins on the cell surface. A number of selection systems may be used, including but not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, *Cell* 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, *Proc. Natl. Acad. Sci. USA* 48:2026), and adenine phosphoribosyl transferase (Lowy *et al.*, 1980, *Cell* 22:817) genes which can be employed in tk<sup>-</sup>, hgprt<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for *dhfr*, which confers resistance to methotrexate (Wigler *et al.*, 1980, *Natl. Acad. Sci. USA* 77:3567; O'Hare *et al.*, 1981, *Proc. Natl. Acad. Sci. USA* 78:1527); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981), *Proc. Natl. Acad. Sci. USA* 78:2072); *neo*, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin *et al.*, 1981, *J. Mol. Biol.* 150:1); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, 1984, *Gene* 30:147). Additional selectable genes have been described, namely *trpB*, which allows cells to utilize indole in place of tryptophan; *hisD*, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, 1988, *Proc. Natl. Acad. Sci. USA* 85:8047); *ODC* (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., 1987, In: *Current Communications in Molecular Biology*, Cold Spring Harbor Laboratory ed.) and glutamine synthetase (Bebbington *et al.*, 1992, *Biotech* 10:169). Alternatively, homologous recombination into an advantageous site may be used to introduce DNA into cells (e.g., murine embryonic stem cells).

When interrogating a tethered ligand or tethered ligand array with a recombinant cell, it has been observed that better signal results when the cells are subcloned ("split") 1-2 days before the assay.

### 5.1.2 Characterization of Orphan Receptors

In one embodiment, the reagents and methods of the invention are useful for identifying ligands that bind to an orphan receptor. There are many putative receptors having various predicted biological functions that have no known ligands. According to the invention, cloned orphan receptors are recombinantly expressed, as described herein, and the binding of the orphan-receptor-expressing cell to a variety of tethered ligands is tested. Based on the profile of binding, a ligand or ligands can be assigned to the receptor.

Usually, one of skill is able to assign an orphan receptor to a specific class of receptors based on sequence homology or other characteristics. It will be recognized that, in such a case, the tethered ligand corresponding to the orphan receptor will usually, or initially, be interrogated with ligands known or believed to bind receptors of this class. For example, an orphan receptor with homology to a CXC chemokine receptor would initially be interrogated with ligands known to interact with CXC chemokine receptors. Similarly, an orphan receptor with homology to one or more classes of chemokine receptors will usually, or initially, be interrogated with chemokine ligands.

## 5.2 Interrogation with populations of cells

In various embodiments, the invention provides methods for analyzing the receptor (e.g., chemokine receptor) profile of a population of cells. The terms "receptor profile" or "profile of receptor expression" (used interchangeably) refer to the complement of cell-surface displayed receptors (i.e., membrane bound extracellular receptors) on a cell type (e.g., cell line or purified cell type) or in a heterogeneous population of cells (e.g., a cell sample from a patient), as detected in an assay. Thus, in an assay for a set of chemokine receptors, the term "receptor profile" for a cell line will refer to the set of those chemokine receptors displayed by cells of the cell line (without regard to other receptors, e.g., growth factor receptors, displayed). Similarly, the term "receptor profile" for a heterogeneous mixture of cells refers to the set receptors displayed by the cells of the population, even though not every cell may display the same set of receptors. As is discussed *infra*, the receptor profile of a cell population can also include information about the number and/or type of cells in the population binding each of several receptors.

Determining the receptor profile for populations of cells is useful in a variety of applications including diagnosis or prognosis of diseases in patients (or nonhuman animals, including medical models); profiling populations of individuals; assessment of the effect of a drug or treatment on a patient, and others.

### 5.2.1 Diagnosis and Prognosis

Different types of cells express different extracellular receptors. Moreover, the receptors expressed by any particular cell type may vary with the developmental stage of the cell, the cell environment, and location. For example,



virally-infected cells may express different receptors than similar uninfected cells. Further, the presence or absence of cells expressing certain receptors (i.e., a cell population with a particular receptor profile) at a particular location is indicative of the state of health of an individual. Examples include the presence of leukocytes and other cells at sites of inflammation, and the presence of malignant cells in a sample.

Accordingly, the present invention provides diagnostic methods in which a population of cells is obtained from a patient suspected of suffering from a disease and the receptor profile for the population is determined using the assay methods described herein (i.e., contacting the cell population with one or more immobilized tethered ligand fusion proteins and identifying the ligands bound by the cells in the population). The receptor profile is then compared to a profile characteristic of the disease state and a profile characteristic of a healthy state, and a diagnosis is based on the comparison.

Examples of populations of cells include, without limitation, (1) cells from disease tissues and fluids, (2) cells from patient populations, (3) cells taken from patients at various times during the progression of disease or treatment. Thus, in one embodiment, cells are obtained from a tissue or fluid from an individual suffering from a disease, e.g., synovial fluid from an individual with rheumatoid arthritis; cerebral-spinal fluid from individuals with multiple sclerosis and Alzheimer's disease, bronchial alveolar lavage (BAL) fluid from patients with asthma, sarcoidosis, tuberculosis, adult respiratory distress syndrome and other inflammatory and infectious diseases, fetal cells, tissues, and biopsy (e.g., from tissues such as muscle, bone marrow, lymph node, liver, brain, and others). The reagents and methods of the invention are used to determine the population of receptors (e.g., chemokine receptors or "CKRs") expressed in the population and compared, for example, to the CKRs expressed in normal (e.g., non-disease) tissue. The normal and nondiseased tissue may be from the same subject, or from two or several different subjects. Differentially expressed CKRs define specific molecular disease targets for therapeutic intervention.

Cells may be obtained from patients suffering from any of a variety of diseases including without limitation diseases and conditions associated with inflammation, infection and cancer, e.g., (1) inflammatory or allergic diseases such as systemic anaphylaxis or hypersensitivity responses, drug allergies, insect sting allergies; inflammatory bowel diseases, such as Crohn's disease, ulcerative colitis,

ileitis and enteritis; vaginitis; psoriasis and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria; vasculitis; spondyloarthropathies; scleroderma; respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, and the like, (2) autoimmune diseases, such as arthritis (rheumatoid and psoriatic), multiple sclerosis, systemic lupus erythematosus, diabetes, glomerulonephritis, and the like, (3) graft rejection (including allograft rejection and graft-vs-host disease), and (4) other diseases in which undesired inflammatory responses are to be inhibited (e.g., atherosclerosis, myositis); cancers, diseases in which angiogenesis or neovascularization play a role (neoplastic diseases, retinopathy and macular degeneration), infectious diseases and immunosuppressive diseases.

Further information about the population is obtained by quantitating and/or characterizing the cells bound to one or more of the ligands (e.g., by counting, sorting, immunostaining, and the like).

#### 5.2.2 Drug Assessment

The reagents and methods of the invention are also useful for monitoring the effect of a drug or treatment on a patient. Cells from an individual or nonhuman animal are obtained from prior to and after treatment for a disease, or before or after onset or progression of a disease, before or after exposure to a pathogen, before and after immunization, and the like, and changes in the profile of receptor expression assayed by comparing the results. Further information about the populations is obtained by quantitating and/or characterizing the cells bound to one or more of the ligands (e.g., by counting, sorting, immunostaining, and the like). The drug assessment can be carried out repeatedly during a course of treatment.

#### 5.2.3 Profiling populations

In another related embodiment, the methods of the invention are used for "profiling" populations of individuals. In an exemplary embodiment, it is useful to determine the receptor profile (e.g., CKR profile) of cells (e.g., blood cells or leukocytes) from specific populations of individuals in order to group the individuals into cohorts of individuals expressing a particular receptor profile, or into cohorts of individuals expressing a particular receptor (e.g., on particular cells). For example, to plan a clinical trial for a CCR3 agonist to inhibit the binding of eotaxin to CCR3

expressed on eosinophils (e.g., for treatment to retard the activity of eosinophils in asthma), it will be useful to profile a large sample of individuals in the general population for the presence of CCR3. A subset of the population deficient in eosinophil binding to eotaxin-stalkokines can be identified and excluded from the clinical trial. Similarly, CCR3 expression can be correlated with other characteristics or markers (e.g., gender, age, or genetic or ethnic variants).

### 5.3 Interrogation with membrane preparations

In addition to interrogating with recombinant or naturally occurring cells expressing the receptors of interest, a membrane preparation may be obtained from such cells and or cell homogenates may be used in the assays of the invention. Membrane preparations may be prepared using routine methods, e.g., hypotonic lysis and centrifugation (see, e.g., Dairaghi et al, 1999, *J. Biol. Chem.* 274:21569-74).

## 6. METHODS OF DETECTION

Cell binding to tethered ligands and arrays can be detected and quantitated in a number of ways. Cells can be observed (and counted) visually (i.e., using a microscopy) with or without prior staining. Alternatively, automated and semiautomated detection systems (e.g., fluorometric imaging plate readers, scintillation counters, "96-well" plate readers) may be used to measure the signal associated with particular tethered ligands (e.g., sites on an array), particularly when the receptor bearing cells are labeled with a detectable label. As used herein, a "detectable label" has the ordinary meaning in the art and refers to an atom (e.g., radionuclide), molecule (e.g., fluorescein), or complex, that is or can be used to detect (e.g., due to a physical or chemical property), indicate the presence of a molecule or to enable binding of another molecule to which it is covalently bound or otherwise associated. Detectable labels suitable for use in the present invention include any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Labels useful in the present invention include biotin for staining with labeled streptavidin conjugate, magnetic beads (e.g., Dynabeads<sup>TM</sup>), fluorescent dyes (e.g., fluorescein, Texas red, rhodamine, green fluorescent protein, enhanced green fluorescent protein, lissamine, phycoerythrin, Cy2, Cy3, Cy3.5, Cy5, Cy5.5, Cy7, FluorX [Amersham],

SyBR Green I & II [Molecular Probes], and the like), radiolabels (e.g.,  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^{32}\text{P}$ ), enzymes (e.g., hydrolases, particularly phosphatases such as alkaline phosphatase, esterases and glycosidases, or oxidoreductases, particularly peroxidases such as horseradish peroxidase, and others commonly used in ELISAs), substrates, cofactors, inhibitors, chemiluminescent groups, chromogenic agents, and colorimetric labels such as colloidal gold or colored glass or plastic (e.g., polystyrene, polypropylene, latex, etc.) beads. Patents teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Means of detecting such labels are well known to those of skill in the art.

In one embodiment, the association of receptor and tethered ligand is detected using the microswitch technology described at <http://www.ece.neu.edu/edsnu/zavracky/mfl/programs/relay/relay.html>.

If desired, cells may be fixed (e.g., using cross-linking agents such as glutaraldehyde) prior to detection.

As noted *supra* it will sometimes be convenient to stain receptor-bearing cells prior to or after contacting the cells with the tethered ligands according to the invention. In one illustrative embodiment, fluorescein isothiocyanate (FITC) labeled anti CD20 antibody is used to mark B cells, and rhodamine labeled anti CD3 is used to mark T cells. Visualization and quantitation under UV illumination and the correct wavelength filter sets would discern between the different cell types.

## 7. DATA ANALYSIS

Modes of analysis of the binding of receptor bearing cells to the tethered ligands and arrays of the invention will be apparent to one of skill following the guidance of this disclosure. Typically, analysis of the binding of the receptor expression cell line to the tethered array is done after quantitation of cell binding by enumeration, fluorescence, or scintillation. Usually, binding is done in triplicate and the average (mean) of the value is used for analysis.

Initially a baseline of non-specific binding is determined. The measurements for the wells in which a non-binding tethered ligand (e.g. a tethered chemokine of a subclass and does not bind a test chemokine receptor) serves as an appropriate baseline. Additional appropriate baselines can be determined by the binding measured for a non-ligand containing immobilized stalk (i.e., a "tethered

ligand" lacking the ligand domain). Additionally, a mock-transfected control containing supernatant of a cell line that was processed through the transfection protocol but does not contain any plasmid capable of making the tethered ligand (such as the parental vector pcDNA3.1) may be used. A positive control also may  
5 be used in some embodiments (e.g., a tethered ligand known to bind to a particular receptor-bearing cell). Suitable positive controls exhibit binding at least about two-fold higher than base line, often about 10-fold higher than base line, and frequently at least about 100-fold higher than base line.

Authentic binding (e.g. binding occurring through the transfected  
10 receptor and the tethered ligand) can be verified by competition assays. In one embodiment, a soluble form of the ligand (i.e., corresponding to the ligand motif of the tethered ligand) or a soluble (non-immobilized) form of the tethered ligand is added in excess (e.g. 100 fold molar excess) to the cell line expressing the receptor for a pre-incubation period (e.g., 5 to 20 minutes, typically 15 minutes). The effect  
15 on binding is determined as described supra. Authentic binding is inhibited by the addition of soluble ligand. Occasionally, the binding can be mediated through endogenous receptors being expressed on the cell line being used. Use of the parental (i.e, non-recombinant) cell line as a control in a parallel analysis is used to detect this.

## 20 8. BINDING AGONISTS AND ANTAGONISTS

In various embodiments, the invention provides a method for identifying the ability of a test compound to modulate the interaction between a ligand and a receptor, between several ligands and a receptor, or between several  
25 ligands and several receptors.

In one embodiment, the binding assays of the invention, using either an array of tethered ligands or one or more individual tethered ligands is carried out in the presence and absence of a test compound. Decreased binding of receptor-bearing cells to the tethered ligand(s) in the presence of the test compound indicates  
30 that the test compound is acting as an antagonist of the interaction. Increased binding of receptor-bearing cells to the tethered ligand(s) in the presence of the test compound indicates that the test compound is acting as an agonist of the interaction. Because ligand receptor interactions have physiological effect (e.g., on cell metabolism), modulators of the interaction are expected to modulate that effect, and

such modulators are often therapeutically useful, e.g., for producing pharmaceutical compositions.

The test compounds referred to *supra* may be any of a large variety of compounds, both naturally occurring and synthetic, organic and inorganic, and including polymers (e.g., oligopeptides, polypeptides, oligonucleotides, and polynucleotides), small molecules, antibodies, sugars, fatty acids, nucleotides and nucleotide analogs, analogs of naturally occurring structures (e.g., peptide mimetics, nucleic acid analogs, and the like), and numerous other compounds. Typically, test agents are added at a variety of concentrations, e.g., ranging from about 1 ng to about 1 g/ml, more often between 1 ug/ml and 1 mg/ml. Typically, test agents are added at a variety of molar concentrations, e.g. ranging from 1 picomolar to 1 molar, more often 1 nanomolar to 1 millimolar. Typically, test agents are added at a variety of concentrations, e.g. ranging from about 1 part per billion to about 1 part per hundred, more often from about 1 part per million to about 1 part per thousand.

In one embodiment, the "test compound" is a soluble form of the ligand moiety of the tethered ligand with which the receptor interacts. For example, the chemokines ELC, TECK and SLC have been demonstrated to inhibit (i.e., by competition) the binding of a CCR10-expressing cell to a tethered ligand having an ELC ligand domain (i.e., an ELC-stalkokine). See Example 1. More often, the "test compound" is an "synthetic test compound" usually a non-polypeptide compound, i.e., not having a sequence of a naturally occurring polypeptide.

In assessing the modulatory activity of a test agent, the test agent may be preincubated with the receptor bearing cell(s) prior to contacting of the cell(s) and the tethered ligand(s); added after the contacting of the cell(s) and the tethered ligand(s); added simultaneously with the contacting of the cell(s) and the tethered ligand(s). Test agents may also be administered to animals or in vitro or ex vivo cells or tissues to assay the effects of binding of populations of cells in a particular target tissue, or animal model.

In one embodiment of the invention, the effects of test agent(s) on the interaction between a particular tethered ligand and a specific receptor are tested. This approach is useful, for example, for validating an apparent interaction between a receptor (e.g., an orphan receptor) and a specific ligand and for high-throughput screening of agents that inhibit a specific receptor-ligand interaction (thereby modulating the biological function of the receptor or ligand). Methods for high-

throughput screening are known (see, e.g., Williams, 2000, *Curr. Opin. Biotechnology* 11:42-46, and references cited therein).

In another embodiment, the arrays of the invention are used to assay the effect of test agents on the binding profile of a population of cells to a plurality of ligands.

## 9. KITS

In one aspect the invention provides kits (e.g., containers) containing (1) a tethered ligand array of the invention, or (2) one or several of the tethered ligand fusion proteins of the invention (e.g., a combination of at least 2, at least 5, or at least 10 different tethered ligands), e.g., in separate vials or (3) one, two or a plurality of the tethered ligand fusion proteins, with at least two of the tethered ligand fusion proteins immobilized different solid substrates (e.g., plates, slides, beads, or the like), e.g., different substrates (such as different beads).

## 10. EXAMPLES

The following examples are offered to illustrate, but not to limit, the claimed invention.

### EXAMPLE 1

This example demonstrates the use of the invention to determine the ligand specificity of an orphan receptor.

#### A. Abbreviations

ELC, EBI1 ligand chemokine; SLC, secondary lymphoid-tissue chemokine; TECK, thymus expressed chemokine; HEK293, human embryonic kidney 293 cells; PEI, polyethylenimine; CCR, CC chemokine receptor.

#### B. Materials and Methods

Human, viral and murine recombinant chemokines were obtained from R&D Systems (Minneapolis, MN; [http://cytokine.rndsystems.com/cyt\\_cat/cyt\\_cat.html](http://cytokine.rndsystems.com/cyt_cat/cyt_cat.html)). <sup>125</sup>I-labeled ELC and TECK were obtained from Amersham.

Full length CCR10 expression constructs were made in pIRESpuro expression vector (Clontech, Palo Alto, CA) with a FLAG epitope tag and prolactin signal sequence, and used to generated stable transfectants in HEK293 cells. Transient and stable transfections for CCR10 and stalkokines were done using Superfect reagent (Qiagen, Valencia, CA) following manufacturer's protocol. Stably-transfected cells were generated by selection in 2 ug/mL puromycin for 7 days, and expression was confirmed by FACS analysis of the FLAG epitope using anti-FLAG M1 (Sigma, St. Louis, MO) and 2' anti-mouse PE conjugate (Coulter Immunotech, Miami, FL).

#### C. Stable expression of CCR10 protein.

The cloning and characterization of a new human chemokine receptor, designated "CCX CKR" or "CCR10" is described in Gosling *et al.*, 2000, "Cutting edge: identification of a novel chemokine receptor that binds dendritic cell- and T cell-active chemokines including ELC, SLC, and TECK" *J Immunol.* 164:2851-6, incorporated herein by reference in its entirety for all purposes. The coding sequence of CCR10 is shown in Fig. 2

To assess the functional properties of the protein encoded by the CCR10 cDNA, including its potential chemokine binding profile, expression plasmids encoding CCR10 plus an added N-terminal Flag epitope were constructed. This allowed for detection and selection, using an anti-Flag mAb, of the most highly expressing stable transfectants. Human embryonic kidney 293 (HEK293) cells stably expressing the M1 flag epitope-tagged CCR10 were confirmed by FACS, and were selected for further analysis (HEK293-CCR10 cells).

#### D. Receptor interrogation by adhesion to stalkokines.

To determine ligand binding to CCR10, HEK293-CCR10 cells were used to interrogate chemokine "stalkokines" (SK), i.e., molecules in which discrete chemokine domains were engineered to be tethered to the end of an extended stalk structure. Stalkokines were interrogated using 8-well chamber slides coated first with anti-His anchoring antibody (10 ug/ml in PBS overnight at RT), which were washed and 'blocked' (2% FBS/0.5% BSA in PBS); treated with 250 ul of HEK293 cell stalkokine supernatants (1 hr at 37C), and incubated with 500,000 HEK293-CCR10 transfectants (1.5 hrs at RT). Inhibition of adhesion by competition with



soluble chemokines was done by incubating cells with 5-10 ug/ml of recombinant chemokines. In all cases, nonadherent cells were removed by washing in PBS; remaining adherent cells were fixed with 1% glutaraldehyde, photoimaged and counted. As a primary screen, this adhesion would reveal putative receptor-ligand interactions.

CCR10-expressing cells adhered very well to ELC stalkokines (ELC-SK; Fig 3A). Furthermore ELC-SK mediated adhesion was abolished in the presence of soluble native ELC as a competitor (Fig 3A, top row). A significant reduction in ELC-SK mediated adhesion of HEK293-CCR10 cells in the presence of soluble SLC, as well as soluble TECK, but not soluble MCP-3 (Fig. 3A, bottom row) was also observed. These experiments were performed and quantitated over several independent trials, an example of which is given in Fig. 3B, and were found to be highly reproducible. Moreover, radiolabeled ELC was used in a traditional homologous competition assay in the presence of increasing concentrations of unlabeled ELC. The results revealed significant binding of ELC to HEK293-CCR10 cells, but not to wildtype (wt) HEK293 cells (Fig 3C). Nearly identical results were obtained in homologous competition of radiolabeled TECK with cold TECK (not shown). Taken together, the stalkokine-based adhesion and radiolabeled ligand binding/homologous competition assays indicate that CCR10 is a new chemokine receptor that bound a novel complement of chemokines.

The spectrum of ligands that bind to CCR10 includes ELC, SLC, and TECK with high affinity, and BLC and vMIPII with lower affinity.

## EXAMPLE 2

This example describes the construction and expression of stalkokines, and the adhesion of receptor-expressing cells to immobilized stalkokines.

### A. Preparation of Stalkokine expression plasmids

#### 1. Amplification of CK Ligand Sequences

Chemokine ("CK") coding sequences were generated using the polymerase chain reaction (PCR) with oligonucleotides synthesized based on the published literature, as shown in Table 5. The amplified CK coding sequence was cloned into an expression vector encoding a human fractalkine stalk region coding

sequence and a c-terminal epitope tag (poly-histidine). Starting material for the PCR reactions included previously cloned plasmids, cDNA libraries, and cDNA generated by reverse transcription of RNA, as indicated in the table.

Amplification was carried out using oligonucleotides containing unique restriction endonuclease sites (EcoR1 in most case, alternatively EcoRV or SmaI) for cloning purposes. The amplified chemokine fragments included the initiation methionine codon and leader peptide sequence, resulting in a secreted product.

For example, a polynucleotide encoding the entire MCP2 chemokine was generated by PCR amplification. The amplification oligonucleotides each included an EcoRI site. The PCR fragment contained the regions encoding the whole, unprocessed chemokine (leader sequence plus mature protein), including the codons for the initiating methionine, leader sequence, and entire chemokine coding sequence up to, but not including the stop codon (amino acid 1 (met) to 109 (pro) inclusive, Genbank accession number X99886).

Following amplification of the chemokine domain sequence, the DNA fragment was subcloned into the EcoRI (or EcoRV site) of the modified vector based on pcDNA3.1, encoding a human Fractalkine mucin stalk sequence (described below). See Figure 4. The ligand motif is designed to be in frame by the construction of the oligonucleotides for the PCR synthesis of the fragment and the sequence of the construct is determined by DNA sequencing.

Table 5

Chemokine	Forward Primer	Reverse Primer	Tissue Source
MDC GenBank accession number: U83171	5'-GGT GAA TTC ATG GCT CGC CTA CAG ACT GC (SEQ ID NO:23)	3'-GGT GAA TTC TTG GCT CAG CTT ATT GAG AAT CA (SEQ ID NO:24)	RT-PCR from human immature dendritic cell RNA
HCC1 GenBank accession number: Z49270	5'-GGT GAA TTC ATG AAG ATC TCC GTG GCT GCC (SEQ ID NO:25)	3'-GGT GAA TTC GTT CTC CTT CAT GTC CTT GAT ATA G (SEQ ID NO:26)	PCR from human placenta cDNA library
SLC GenBank accession number: AF001979, AB002409	5'-GGT GAA TTC ATG GCT CAG TCA CTG GCT CTG (SEQ ID NO:27)	3'-GGT GAA TTC TGG CCC TTT AGG GGT CTG TG (SEQ ID NO:28)	PCR from human EST clone
MIG GenBank Accession number NM 002416	5' GGT GAA TTC ATG AAG AAAA GT GGT GTT CTT TTC C (SEQ ID NO:29)	3' GGT GAA TTC TGT AGT CTT CTT TTG ACG AGA ACG (SEQ ID NO:30)	reverse transcription of human fetal brain mRNA
IP10 GenBank	5' GGT GAA TTC ATG AAT CAA ACT GCG ATT	3' GGT GAA TTC AGG AGA TCT TTT AGA CAT	human cDNA pool

Accession number NM 001565	CTG A (SEQ ID NO:31)	TTC CTT G (SEQ ID NO:32)	
H174 / I-TAC GenBank accession number AF002985	5' GTG GAA TTC ATG AGT GTG AAG GGC ATG GC (SEQ ID NO:33)	3' GGT GAA TTC AAA ATT CTT TCT TTC AAC TTT TTT GA (SEQ ID NO:34)	human placental cDNA library
Eotaxin1 GenBank accession number U46573	5' GGT GAA TTC ATG AAG GTC TCC GCA GCA CTT C (SEQ ID NO:35)	3' GGT GAA TTC TGG CTT TGG AGT TGG AGA TTT TTG (SEQ ID NO:36)	human EST
TARC GenBank accession number NM 002987	5' GGT GAA TTC ATG GCC CCA CTG AAG ATG CT (SEQ ID NO:37)	3' GGT GAA TTC AGA CCT CTC AAG GCT TTG CAG (SEQ ID NO:38)	reverse transcription of immature dendritic cell mRNA
MIP3alpha GenBank accession number U77035	5' GGT GAA TTC ATG TGC TGT ACC AAG AGT TTG C (SEQ ID NO:39)	3' GGT GAA TTC CAT GTT CTT GAC TTT TTT ACT GAG G (SEQ ID NO:40)	human EST
ELC GenBank accession number U77180, NM 006274	5' GGT GAA TTC ATG GCC CTG CTA CTG GCC CT (SEQ ID NO:41)	3' GGT GAA TTC ACT GCT GCG GCG CTT CAT CT (SEQ ID NO:42)	human EST
TECK GenBank accession number NM 005624	5' GGT GAA TTC ATG AAC CTG TGG CTC CTG GC (SEQ ID NO:43)	3' GGT GAA TTC CAG TCC TGA ATT AGC TGA TAT CAG (SEQ ID NO:44)	human placental cDNA library
PARC GenBank accession number AB012113	5' GGT GAA TTC ATG AAG GGC CTT GCA GCT GC (SEQ ID NO:45)	3' GGT GAA TTC GGC ATT CAG CTT CAG GTC GC (SEQ ID NO:46)	human placental cDNA library
SLC GenBank accession number AF001979	5' GGT GAA TTC ATG GCT CAG TCA CTG GCT CTG (SEQ ID NO:47)	3' GGT GAA TTC TGG CCC TTT AGG GGT CTG TG (SEQ ID NO:48)	human EST
I309 GenBank accession number AA931884	5' GGT GAA TTC ATG CAG ATC ATC ACC ACA GCC C (SEQ ID NO:49)	3' GGT GAA TTC TTT TCT TTT TGA CGG GCA GTG C (SEQ ID NO:50)	human EST
MIP1alpha GenBank accession number AA031820	5' GGT GAA TTC ATG CAG GTC TCC ACT GCT GC (SEQ ID NO:51)	3' GGT GAA TTC GGC ACT CAG CTC TAG GTC GCT (SEQ ID NO:52)	human cDNA pool
MCP-4 GenBank accession number U59808	5' GGT GAA TTC ATG AAA GTC TCT GCA GTG CTT CTG (SEQ ID NO:53)	3' GGT GAA TTC AGT CTT CAG GGT GTG AGC TTT CC (SEQ ID NO:54)	reverse transcription of human brain mRNA
MCP1 GenBank accession number AA024753	5' GGT GAA TTC ATG AAA GTC TCT GCC GCC CTT (SEQ ID NO:55)	3' GGT GAA TTC AGT CTT CGG AGT TTG GGT TTG C (SEQ ID NO:56)	human EST
MCP2 GenBank accession number NM005623	5' GGT GAA TTC ATG AAG GTT TCT GCA GCG CT (SEQ ID NO:57)	3' GGT GAA TTC TGG CTT CAG ATT TTG AAA TAT TTG (SEQ ID NO:58)	reverse transcription of immature dendritic cell mRNA
SDF1a GenBank accession number L36034	5' GGT GAATTC ATG AAC GCC AAG GTC GTG G (SEQ ID NO:59)	3' GGT GAA TTC CTT GTT TAA AGC TTT CTC CAG GT (SEQ ID NO:60)	human EST

MCP5	5' GGT GAT ATC ATG AAG ATT TCC ACA CTT CTA TGC C (SEQ ID NO:61)	3' GGT GAT ATC GCC TAG ACA TGA AGG TTC AAG GAT G (SEQ ID NO:62)	
HCC2	5' GGT GAA TTC ATG AAG GTC TCC GTG GCT GC (SEQ ID NO:63)	3' GGT GAA TTC TAT TGA GTA GGG CTT CAG CTT T (SEQ ID NO:64)	
CLONE 391	5' GGT GAA TTC ATG AAG GTC TTC TCC TTG GTC ATG (SEQ ID NO:65)	3' GGT GAA TTC CGT TGA GGT GTT GCT CAG CTT C (SEQ ID NO:66)	
BCA1	5' GGT GAT ATC ATG AAG TC ATC TCG ACA TCT CTG (SEQ ID NO:67)	3' GGT GAT ATC GGG AAT CTT TCT CTT AAA CAC TGG (SEQ ID NO:68)	
PF4a	5' GGT GAA TTC ATG AGC TCC GCAGCC GGG TTC (SEQ ID NO:69)	3' GGT GAA TTC ACT CTC CAA AAG TTT CTT AAT TAT TTT (SEQ ID NO:70)	
PBP-like1	5' GGT GAA TTC ATG CCA CCC TGC AGC TGT G (SEQ ID NO:71)	3' GGT GAA TTC TAA AGC CAT TGT GAA TAT GAT CTG (SEQ ID NO:72)	
EOTAXIN 2	5' GGT CCC GGG ATG GCA GGC CTG ATG ACC AT (SEQ ID NO:73)	3' GGT CCC GGG GCA GGT GGT TTG GTT GCC AG (SEQ ID NO:74)	
MIP1beta	5' GGT GAA TCC ATG AAG CTC TGC GTG ACT GTC C (SEQ ID NO:75)	3' GGT GAA TTC GTT CAG TTC CAG GTC ATA CAC GTA (SEQ ID NO:76)	
MIP-3	5' GGT GAA TTC ATG AAG GTC TCC GTG GCT GC (SEQ ID NO:77)	3' GGT GAA TTC ATT CTT CCT GGT CTT GAT CCG T (SEQ ID NO:78)	
MRP-1	5' GGT GAA TTC ATG AGA AAC TCC AAG ACT GCC A (SEQ ID NO:79)	3' GGT GAA TTC AGC AAT GAC CTT GTT CCC AGA T (SEQ ID NO:80)	
Lymphotactin (LYNT)	5' GGT GAA TTC ATG ATA CTT CTC ATC CTG GCC C (SEQ ID NO:81)	3' GGT GAA TTC GCC AGT CAG GGT CAC AGC TG (SEQ ID NO:82)	
HCC4	5' GGT GAA TTC ATG AAG GTC TCC GAG GCT GC (SEQ ID NO:83)	3' GGT GAA TTC CTG GGA GTT GAG GAG CTG GG (SEQ ID NO:84)	
MIP-1gamma	5' GGT GAA TTC ATG AAG CCT TTT CAT ACT GCC C (SEQ ID NO:85)	3' GGT GAA TTC TTG TTT GTA GGT CCG TGG TTG T (SEQ ID NO:86)	
IL-8	5' GGT GAT ATC ATG ACT TCC AAG CTG GCC G (SEQ ID NO:87)	3' GGT GAT ATC TGA ATT CTC AGC CCT CTT CAA A (SEQ ID NO:88)	

ENA-78	5' GGT GAA TTC ATG AGC CTC CTG TCC AGC CG (SEQ ID NO:89)	3' GGT GAA TTC GTT TTC CTT GTT TCC ACC GTC C (SEQ ID NO:90)	
GROgamma	5' GGT GAT ATC ATG GCC CAC GCC ACG CTC TC (SEQ ID NO:91)	3' GGT GAT ATC GTT GGT GCT CCC CTT GTT CAG (SEQ ID NO:92)	
GRObeta	5' GGT GAT ATC ATG GCC CGC GCC ACG CTC TC (SEQ ID NO:93)	3' GGT GAT ATC GTT GGA TTT GCC ATT TTT CAG C (SEQ ID NO:94)	
NAP-2	5' GGT GAT ATC ATG AGC CTC AGA CTT GAT ACC ACC (SEQ ID NO:95)	3' GGT GAT ATC ATC AGC AGA TTC ATC ACC TGC C (SEQ ID NO:96)	
GROalpha	5' GGT GAT ATC ATG GCC CGC GCT GCT CTC TC (SEQ ID NO:97)	3' GGT GAT ATC GTT GGA TTT GTC ACT GTT CAG CAT C (SEQ ID NO:98)	
vMIP	5' GGT GAA TTC ATG GCC CCC GTC CAC GTT TT (SEQ ID NO:99)	3' GGT GAA TTC ATG GAC ACC AAG GGC ATC CT (SEQ ID NO:100)	

### B. Stalk Expression Cassette

CK ligand coding sequences were cloned into the EcoR1 or EcoRV sites of the polylinker region of the expression vector pcDNA-FRAC. This vector includes a polylinker upstream of a human fractalkine stalk coding region fused to the myc epitope and poly-histidine coding region used for antibody tethering and subsequent purification. The coding region is under control of a CMV promoter.

pcDNA-FRAC was constructed by inserting a fractalkine mucin stalk coding sequence (corresponding to amino acids 100 to 336 (inclusive) of human fractalkine (Genebank Accession number U84487)) into the EcoRI and HindIII restrictions sites of vector pcDNA3.1(-)/Myc-His version B (Invitrogen) [hereinafter, "pcDNA3.1"]. The resulting vector (pcDNA3.1-FRAC) was used for expression of various chemokine ligands. This was accomplished by inserting a PCR-amplified ligand sequence into the vector upstream of, and in frame with, the stalk domain. The resulting vectors (and the encoded proteins) are generally referred to as "chemokine'-FRAC", e.g., "MDC-FRAC", "TARC-FRAC" and the like.

A region of the multiple cloning site polylinker region from pcDNA3.1 is present in between the last amino acid of the chemokine ligand domain and the first amino acid (glycine) of the fractalkine mucin stalk region. Insertion of the PCR fragment encoding the chemokine into the polylinker puts the coding regions in frame, resulting in the proper full length translation of the fusion protein. The

polylinker region encodes various amino acids, depending on the restriction sites used for the cloning of the chemokine domain. For example, if the EcoR1 linkered chemokine domain is cloned into the EcoR1 site of the pcDNA3.1-FRAC, the linker region will encode the additional amino acids glutamate-phenylalanine. As another example, if the EcoRV linkered chemokine domain is cloned into the EcoRV site, the linker region will encode the additional amino acids alanine-glutamate-phenylalanine. As yet another example, if the SmaI linkered chemokine domain is cloned into the EcoRV domain (as a blunt end ligation), the linker region will encode the additional amino acids alanine-isoleucine-proline-alanine-glutamate-phenylalanine.

The final plasmid is prepared in bulk using standard protocols and the Qiagen purification system to generate pure plasmid in milligram amounts. Plasmid was stored in 10 mM Tris, 1 mM EDTA at 20°C.

#### C. Expression of Stalkokine in cell culture

The stalkokine synthesis is generated in *in vitro* cell culture using HEK293 (ATCC No. CRL-1573) or 293T (Pear et al, 1993, *PNAS* 80:8392-6) cell lines transiently transfected with the stalkokine expression plasmid. These cell lines were grown in DMEM supplemented with 10% FBS containing antibiotics, at a 60% confluence prior to transfection. On the day of transfection, the cells were transfected by the following protocol:

The cells were washed with PBS. 17 ug of plasmid DNA was resuspended in 825 ul of "Opti-MEM" reduced serum medium (Life Technologies, Rockville, MD)). Superfect reagent (Qiagen) was added to the plasmid, mixed, and allowed to incubate at room temperature for 20 minutes. It is then added to the cells and incubated at 37°C, 5% CO<sub>2</sub> for 3 hours. Next, the cells are washed with PBS, and complete DMEM media containing 10% FBS and antibiotics are added. After overnight culture, the media was aspirated and replaced with DMEM with 2.5% FBS and antibiotics.

The supernatants were harvested from the transfected cells at 24, 48, 72. The transfected culture was supplemented with fresh media and incubated as above until the next harvest time. The supernatant was centrifuged to remove any cellular debris, and then transferred to a fresh tube. Several aliquots are made and the samples are stored at -20°C until required. In some experiments, protease inhibitor cocktails were added to the clarified supernatant prior to storage.

Purification of the stalkokines was carried out using poly histidine affinity chromatography. Large scale transfections were done and the supernatants after culture were dialyzed to remove any very small molecule contaminants. The supernatant was passed over the nickel column twice, eluted with imidazole, and the eluate was dialyzed into PBS and analyzed. Usually, significant purification was achieved with reasonable recoveries. This material was compared with the unpurified stalkokine by Western analysis, and total protein was determined. The material purified was typically 50-75% pure.

Purification is useful for a number of reasons: it removes contaminant molecules that could have a detrimental effect on stability of the stalkokine, allows easy comparison of equivalent amounts of material in assays, and facilitates quality control in preparing tethered ligand arrays.

#### D. Monitoring Expression By Western Blotting

The fusion protein is analyzed by Western blot analysis to confirm the expected length of the fusion protein and the presence of the poly histidine tail (which confirms the carboxy terminal is in the correct reading frame). In many cases, the amino-terminal chemokine domain is also interrogated by Western blot analysis looking with chemokine-specific antibodies which will only react with protein made in the correct frame.

The success of the cell transfection was monitored using Western blotting procedures. Typically, 10 ul of supernatant was mixed with a denaturation solution containing reducing agents (DTT) and heated to 100°C for 5 minutes. The protein sample was then loaded onto a 10-20% acrylamide Tricine gel and electrophoresed for 45 minutes at 100 V. The contents of the gel were transferred to a nitrocellulose membrane using standard transfer protocols. Transfer efficiency was determined by completeness of transfer of the prestained molecular weight markers. The membrane was blocked in a solution of Tris buffered saline, BSA and Tween-20. The membrane was then incubated for 1 hour with a primary antibody such as the anti-HIS antibody, or an antibody recognizing the chemokine domain. The primary antibody was washed off with three consecutive washes in TBS for 5 minutes with gentle agitation. The membrane was then incubated for 1 hour with the secondary antibody recognizing the Fc domain of the primary antibody. This secondary antibody has a covalently linked horseradish peroxidase enzyme

attached for subsequent visualization. After washing off the secondary antibody, the membrane was visualized by incubation with a chromagenic substrate DAB until visible bands were generated, then stopped by washing in water. See Figure 5.

#### 5 E. Stalkokine Activity

The functional activity of the chemokine motif of an expressed stalkokine fusion protein was assessed by two methods: chemotaxis and displacement of labeled chemokines.

Fig. 6 shows a displacement assay showing competition by a stalkokine with binding of a radiolabeled tracer chemokine to its cognate receptor. CEM cells expressing CCR4 were used in a binding assay with radiolabeled MDC or TARC, as described in Dairaghi et al, 1999, *J. Biol. Chem.* 274:21569-74.

Briefly, CEM cells were added to radiolabeled chemokine "tracer" (MDC or TARC) in the presence of competitor molecules such as supernatants expressing the soluble MDC-FRAC (a stalkokine with a fractalkine stalk and an MDC ligand domain) or TARC-FRAC (a stalkokine with a fractalkine stalk and an TARC ligand domain). The cells and tracer were incubated and harvested by filtration, capturing the bound tracer. The material was then counted. "Total" is the amount of tracer bound with no competition. "MDC-SK" is the amount of binding in the presence of soluble MDC-FRAC fusion protein, "TARC-SK" is the amount of binding in the presence of soluble TARC-FRAC fusion protein, "ConSup" refers to competition with the control supernatant (conditioned media from CEM cells not expressing any stalkokine), "MDC-CK" is the competition with soluble MDC chemokine, and demonstrated the background binding.

As is evident, there is considerable inhibition of binding of either MDC or TARC tracer to the CEM cells in the presence of the experimental supernatants. Part of this inhibition is due to other elements present in the control sup, but there is a significant further decrease specific to the recombinant MDC-FRAC or TARC-FRAC. The MDC-CK inhibition demonstrates the background binding and the maximal level of competition that one can achieve.

Chemotaxis assays were done on cell types known to possess the receptor for the chemokine ligand (Bacon et al, 1988, *Br J Pharmacol* 95:966-74).



Fig. 7 shows an assay in which CEM cells were "loaded" with INDO-1AM (Molecular Probes, Eugene, OR) and subjected to calcium flux analysis as described in Dairaghi et al, 1997, *J. Biol. Chem.* 272:28206-209.

5 Briefly, CEM cells that express CCR4 were placed in a fluorimeter and monitored over time for changes in fluorescence. Two natural ligands for CCR4 are MDC and TARC. TARC-Fractalkine or MDC-Fractalkine (fusion protein supernatants), or control supernatants, were added at the following times: experiment 1, control supernatant added at 50 sec, TARK-FRAC at 80 sec; experiment 2, control supernatant at 50 sec and MDC-FRAC at 80 sec. The control  
10 supernatant did not result in any calcium influx, where as the TARC-FRAC or MDC-FRAC resulted in modest influx. This experiment demonstrates that the chemokine ligand fusion proteins are capable of causing signaling.

#### F. Adhesion Of Stalkokine To Solid Surfaces

15 Stalkokines were immobilized on glass and polystyrene surfaces by binding to anti-histidine mAbs. The glass substrates were in the form of slides, either 8 well chamber slides, or glass slides printed with hydrophobic Teflon mask to generate wells. Standard microscope slides sizes had between 8 and 30 wells per slide.

20 In one set of experiments, five 8 well chamber slides (Lab-Tek II, Nalge Nunc International, Rochester, NY), 250ul of 10ug/ml anti-his antibody (R&D Systems, Minneapolis, MN) in PBS was added to each well and incubated at room temperature overnight (typically 16 hours). Following, the liquid in each well was aspirated and washed once for approximately 5 minutes with 500ul of PBS. The  
25 wells were then "blocked" (blocked for any additional protein binding sites) with the addition of 250ul of blocking solution (2% fetal bovine serum, 0.5% BSA in PBS) and incubation for 1 hour at room temp (e.g., 20°C). The liquid in each well was aspirated and 250ul of supernatant containing the chemokine-stalk-immobilization domain fusion proteins (SDF1a-FRAC, finetaxin-FRAC, CCL27-FRAC, I309-FRAC,  
30 MIP3a-FRAC, ELC-FRAC, SLC-FRAC, MDC-FRAC, TARC-FRAC, or BRAK-FRAC) or the control supernatant was added to two wells. Each slide contained two wells with control supernatant (from mock-transfected cells) and two wells with the positive control SDF1a (which binds the ubiquitous chemokine receptor CXCR4). The four other wells were for experimental samples. The supernatants were incubated for

one hour at room temperature (typically 20°C), then incubated for an additional hour at 37°C. The supernatants were aspirated off, and each well was washed once for 5 minutes with 500ul of PBS. The slide was then ready for addition of cells for assay.

In a second set of experiments using two 96 well large format glass slides, 50 ul of 10 ug/ml anti-his antibody (R&D Systems, Minneapolis, MN) in PBS was added to each well and incubated at room temperature overnight (typically 16 hours) in a humidified chamber. Following, the liquid in each well was aspirated and washed once for approximately 5 minutes with 50 ul of PBS. The wells were then "blocked" (blocked for any additional protein binding sites) with the addition of 50ul of blocking solution (2% fetal bovine serum, 0.5% BSA in PBS) and incubation for 1 hour at room temp (20°C) in a humidified chamber. The liquid in each well was aspirated and 50ul of supernatant containing one of the 32 chemokine-stalk-immobilization domain fusion proteins shown in Figure 5 was added to 3 wells. Each slide contained 3 wells with the control supernatant (from mock-transfected cells) and 3 wells with the positive control SDF1a (which binds the ubiquitous chemokine receptor CXCR4). The other wells were for experimental samples. The supernatants were incubated for one hour at room temperature (typically 20°C) in a humidified chamber, then incubated for an additional hour at 37C again in a humidified chamber. The supernatants were aspirated off, and each well was washed once for 5 minutes with 50 ul of PBS. The slide was then ready for addition of cells for assay.

#### G. Expression Of Chemokine Receptors In Cell Culture

Several types of cells were used in binding experiments to stalkokines of the invention, including transfected mouse myeloma NSO cells containing a mammalian expression vector driving expression of the receptor of interest, e.g., CCR1, CCR4, CCR6, CCR7, CCR8, CXCR4, CMV US28 (gift of R&D Systems (Minneapolis, MN)). The results of this experiment are summarized in Table 6.

Table 6

Cell types	Some Known Ligands	Binding to Stalkokine form
CCR1-NSO	Ckb8-1, MCP3	no, yes
CCR4-NSO	MDC, TARC	yes, yes
CCR6-NSO	MIP3a	yes

CCR7-NSO	ELC, SLC	yes, no
CCR8-NSO	I309, vMIP1	yes, yes
CCR10-NSO	ELC	yes
CXCR4-NSO	FK, vMIPII	yes

5

\*Negative controls (non-binding stalkokines) were done in every case Ckb8-1 (also MPIF1), ELC (same as MIP3beta), SLC (same as 6Ckine). FK= fractalkine

10

#### H. Contacting Receptor-Expressing Cells and Stalkokine Arrays

In an experiment, CCR1-NSO cells were grown, harvested, and resuspended in PBS at a  $2 \times 10^6$  cells/ml. 250 ul of this cell suspension was added to each well of a 8 well chamber slide, with two wells coated with each of CKb8-1  
15 FRAC, MCP3-FRAC, SDF1a-FRAC (positive control), or mock transfected supernatant (negative control) as previously described. The slides were incubated for 1.5 h at room temp, then the supernatant gently aspirated. The chamber portion of the slide was removed (using the apparatus provided by the supplier) and the slide was washed by gentle immersion into a petri dish containing PBS to a depth of  
20 1cm. This was repeated 3 times in total, and the slide was immersed in a 1% glutaraldehyde solution (in PBS) for 5 minutes at room temp. The cells were inspected and enumerated visually by microscopy. The cells bound to MCP3 but not CKb8 tethered ligands.

#### I. Displacement and competition analysis

In an experiment, CCR4-NSO cells were used to test binding competition. Harvested CCR4 NSO cells were resuspended at  $2 \times 10^6$  cells per ml in PBS. 500 ul aliquots of cells were transferred into eppendorf tubes, and 5 ug of recombinant human chemokine (MDC or TARC, R&D Systems, Minneapolis, MN)  
30 was added (final concentration was 10 ug/ml), mixed briefly, and incubated at room temperature for 10 minutes. Following, 250 ul of the cells were added to each two wells of an 8 well chamber slide that had been precoated with the MDC-FRAC or TARC-FRAC as previously described. 250 ul of cells that were not pre-treated with soluble chemokine were added to additional wells. The slides were incubated for 1.5  
35 hours at room temp, then the supernatant gently aspirated. The chamber portion of the slide was removed (using the apparatus provided by the supplier) and the slide

was washed by gentle immersion into a petri dish containing PBS to a depth of 1cm. This was repeated 3 times in total, and the slide was immersed in a 1% glutaraldehyde solution (in PBS) for 5 minutes at room temperature. The cells were inspected and enumerated visually by microscopy. The cells bound to both the MDC-FRAC and TARC-FRAC, as expected. Competition with either soluble MDC or TARC reduced binding to background levels in both cases, as expected.

In another experiment, small molecule chemical compounds were tested on the binding of vMIP2-FRAC and US28-NSO. Three small organic compounds previously identified as inhibitors of binding between chemokine receptor US28 and vMIP2, were also observed to compete and inhibit binding between vMIP2-FRAC and US28-expressing NSO cells using the present invention.

### EXAMPLE 3

This example describes the binding of a chemokine receptor to an array of stalkokines, to determine the binding specificity of the receptor.

To interrogate the binding of CCR10 to a panel of chemokine elements, an array of tethered ligands is prepared. pc-FRAC-based stalkokines are prepared using at least about 3, at least about 5, at least about 10, at least about 15, at least about 20 or all of the ligand domains listed in Table 5. An array of stalkokines is prepared on a pair of 96 well slides precoated with the anti-polyhistidine antibody overnight and blocked for 1 hour as described *supra*. Each stalkokine is placed on three wells in a 30 ul volume per well and incubated for 1.5 hours in a humidified chamber at 37°C. Also included are negative controls, in triplicate, which include (1) a mock transfected supernatant, (2) an immobilized stalk region (with no chemokine domain), and (3) PBS. A SDF1a stalkokine (which binds CXCR4, a ubiquitously expressed chemokine receptor) is used as a positive control. Each set of positive and negative controls is included on each plate.

The supernatants are aspirated and the wells are washed twice with PBS. The CCR10 cell line is prepared during incubation with the stalkokine. The cell line is trypsinized from the culture flasks, and washed twice with PBS. The cells are resuspended at a concentration of 1 million cells per ml of PBS, 30 ul of the cell suspension are added to each well of plate, and incubated at room temperature without agitation for 1.5 hours at room temperature. The droplets are partially aspirated off (leaving approximately 5 ul, taking care not to disturb the cells resting

on the surface) and placed slowly into a tray with PBS (approximately 150 mls, depth of 1cm). Each of the four edges of the 96 well plate is lifted slowly to the surface of the liquid in each of three consecutive wells. The bound cells are stained with SybrGreen 1 (Molecular Probes, Eugene, OR) for 30 min at a stain diluted 1 to 5000 in PBS. The 96 well plate is removed, drained of excessive liquid, and counted in a Packard Fluorescent plate reader with 480 nm excitation, 530 nm emission filter sets. The results are transferred to an Excel file for further analysis.

\*\*\*

10 All references cited herein are incorporated herein by reference in their entirety and for all purposes to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

15 Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

## WHAT IS CLAIMED IS:

1. An assay device comprising a plurality of different immobilized tethered ligand fusion proteins organized in an array, wherein said ligand-stalk fusion proteins comprise:

- 5           (a) a ligand domain, and  
          (b) a stalk domain.

2. An assay device comprising a plurality of different immobilized tethered ligand fusion proteins organized in an array, wherein said ligand-stalk fusion proteins  
10 comprise:

- (a) a ligand domain;  
          (b) an intermediate stalk domain; and,  
          (c) an immobilization domain,

          wherein said ligand domain and said stalk domain are not associated in a  
15 naturally occurring protein.

3. The device of claim 2 wherein the immobilized tethered ligand fusion proteins comprise a mucin-derived stalk sequence.

20 4. The device of claim 2 wherein the immobilized tethered ligand fusion proteins comprise a fractalkine mucin repeat region sequence.

5. The device of claim 2 wherein at least some of the of immobilized tethered ligand fusion proteins comprise a ligand domain encoding a chemokine.

25

6. A method for identifying an interaction between a receptor and a ligand, said method comprising:

- (a) contacting a cell expressing a receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein; and,  
30           (b) detecting binding of said cell and the fusion protein,  
          wherein binding of the cell to the fusion protein is correlated with an interaction between the receptor and the ligand corresponding to the ligand domain of the fusion protein.

7. The method of claim 6, wherein the tethered ligand is immobilized on an assay device of claim 1.

8. The method of claim 6 wherein the cell expresses a recombinant receptor.

5

9. The method of claim 6 wherein the cell expresses a seven-transmembrane receptor.

10. The method of claim 9 wherein the cell expresses a chemokine receptor or a cytokine receptor.

10

11. The method of claim 6 wherein the cell expresses a orphan receptor.

12. The method of claim 6 wherein an interaction between a receptor and more than one ligand is detected.

15

13. A method for identifying a modulator of an interaction between a receptor and a ligand comprising

20

(a) contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the absence of a test compound, and measuring binding of the cells to the fusion protein;

25

(b) contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the presence of a test compound, and measuring binding of the cells to the fusion protein; and,

30

(c) comparing the level of binding in (a) and (b), wherein decreased binding of cells in the presence of the test compound indicates that the test compound is an inhibitor of the interaction between the receptor and the ligand corresponding the ligand domain of said fusion protein, and wherein increased binding of cells in the presence of the test compound indicates that the test compound is an enhancer of the interaction between the receptor and the receptor and the ligand corresponding the ligand domain of said fusion protein.

14. The method of claim 13 wherein the tethered ligand is immobilized on an assay device of claim 1.

5 15. The method of claim 13 wherein the receptor is a chemokine receptor.

16. A method for detecting a profile of receptor expression in cells in a population of cells comprising:

- 10 (a) contacting said population of cells with an immobilized tethered ligand fusion protein, and
- (b) detecting binding of cells of said population to the fusion protein, wherein binding of a cell in the cell population to the fusion protein is correlated with expression of a receptor that binds the ligand corresponding to the ligand domain of the fusion protein.

15

17. The method of claim 16 wherein step (a) comprises contacting the population with a plurality of different fusion proteins and step (b) comprises detecting binding of cells to zero, one or more than one fusion protein.

20 18. The method of claim 17 comprising detecting binding of cells of said population to zero, one, or more than one tethered ligand fusion protein.

19. The method of claim 17 wherein the tethered ligand is immobilized on an assay device of claim 1.

25

20. The method of claim 16 wherein the population is heterogeneous.

21. The method of claim 20 wherein the population is obtained from synovial fluid, cerebral-spinal fluid, bronchial alveolar lavage (BAL) fluid, or blood.

30

22. The method of claim 16 further comprising quantitating the level of binding to each tethered ligand fusion protein in the array.



23. The method of claim 16 further comprising characterizing the cells bound at each sector of the array.

24. A method for diagnosis comprising

- 5 (a) obtaining a population of cells from a patient suspected of suffering from a disease
- (b) determining a receptor profile for the population; and,
- (c) comparing said receptor profile with a profile characteristic of the disease state.

10

25. A method for diagnosis comprising

- (a) obtaining a population of cells from a patient suspected of suffering from a disease
- (b) determining a receptor profile for the population by
- 15 (i) contacting said population of cells with a device of claim 1; and,
- (ii) identifying a subset of arrayed tethered ligand fusion proteins bound by cells of said population thereby identifying a receptor profile;
- (c) comparing said receptor profile with a profile characteristic of the disease state.

20

26. The method of claim 25, where said determining a receptor profile further comprises characterizing the cells bound at each sector of the array.

27. The method of claim 26 wherein the characterizing is achieved by immunostaining.

25

28. The method of claim 25 further comprising quantitating the binding of cells at each sector of the array.

30 29. The method of claim 24 wherein the disease is an inflammatory or allergic disease, or an autoimmune disease.

30. The method of claim 24 wherein the population is obtained from synovial fluid, cerebral-spinal fluid, bronchial alveolar lavage (BAL) fluid, or blood.

31. A method for detecting an effect of a drug or treatment on a patient comprising:

(a) determining the receptor profile of a population of cells from the patient  
5 for the first time;

(b) administering the drug or treatment to the patient;

(c) determining the receptor profile of a population of cells from the patient  
for a second time,

(d) comparing the receptor profiles in (a) and (c) to determine the effect of  
10 the drug or treatment on receptor-expressing cells in the patient.

32. A method for detecting an effect of a drug or treatment on a patient comprising:

(a) determining the receptor profile of a population of cells from the patient  
15 for the first time, by

(i) contacting said population of cells with a device of claim 1; and,

(ii) identifying a subset of arrayed tethered ligand fusion proteins  
bound by cells of said population thereby identifying a receptor profile;

(b) administering the drug or treatment to the patient;

(c) determining the receptor profile of a population of cells from the patient  
20 for a second time,

(d) comparing the receptor profiles observed in (a) and (c) to determine  
the effect of the drug or treatment receptor-expressing cells in the patient.

25 33. The method of claim 32 further comprising quantitating the level of binding to  
each tethered ligand fusion protein in the array.

34. The method of claim 32 further comprising characterizing the cells bound at  
each sector of the array.

30

35. The method of claim 34 wherein the characterizing is achieved by  
immunostaining.

36. A method for identifying a modulator of an interaction between a receptor and a ligand comprising

(a) contacting cells expressing the receptor or a ligand-binding portion thereof with an immobilized tethered ligand fusion protein in the presence of a test compound and determining the level of binding to the fusion protein,

(b) contacting cells expressing the receptor or a ligand-binding portion thereof with the immobilized tethered ligand fusion protein in the absence of a test compound; and determining the level of binding to the fusion protein, and

(c) comparing the level of binding in (a) and (b) for the immobilized tethered ligand fusion protein,

wherein said ligand-stalk fusion proteins comprise:

(i) a ligand domain;

(ii) an intermediate stalk domain; and,

(iii) an immobilization domain, and

wherein said ligand domain and said stalk domain are not associated in a naturally occurring protein, and,

wherein decreased binding of cells in the presence of the test compound indicates that the test compound is an inhibitor of the interaction between the receptor and the ligand corresponding the ligand domain of said fusion protein, the and wherein increased binding of cells in the presence of the test compound indicates that the test compound is an enhancer of the interaction between the receptor and the receptor and the ligand corresponding the ligand domain of said fusion protein.

37. The method of claim 31 wherein the stalk domain is carboxy-terminal to the ligand domain, and the immobilization domain is carboxy-terminal to the stalk domain.

38. The method of claim 31 wherein the cell expresses a recombinant receptor.

39. The method of claim 31 wherein the cell expresses a chemokine receptor.

40. The method of claim 38 wherein the cell expresses a orphan receptor.

41. The method of claim 36 wherein an interaction between a receptor and more than one ligand is detected.

5 42. A kit comprising at least two different immobilized ligand fusion proteins, wherein said at least two different immobilized ligand fusion proteins are not immobilized on the same substrate.

43. The kit of claim 42 wherein the fusion proteins are immobilized on beads or slides.

10

44. A tethered ligand fusion protein, wherein said ligand domain and said stalk domain are not associated in a naturally occurring protein.

15 45. The tethered ligand fusion protein of claim 44, wherein the ligand domain is other than a chemokine sequence.

1/6




FIG. 1




FIG. 3A

2 / 6

## FIG. 2

(SEQ. ID NO: 1 &amp; 2)

10	20	30	40	50	60	70	80	90
1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890
ATGCTTTGG AACAGACCA GTACACAGT TTTATATAG AGGAATGCA ATGATGTC ACTATGACT ACATGACATA TGAATCATC	M A L E Q N Q S T D Y Y Y B E N E M N G T Y D Y S Q Y E L I							90
270	180							
TGTATCAAG AACATGTCAG AGAATTGCA AAAGTTTCC TCCCTGATT COTACATA GTTTGTC TTTGACTCC AGCAATTC	C I K B D V R E F A K V F L P V F L T I V P V I G L A G N S							180
270								
ATGGTAGTG CATTTAGC CTATTCAAG AACACAGCA CCAACAGCA TGTGTACATC CTGAATTGG CTGTACGCA TTTACTCTT	M V V A I Y A Y Y K K Q R T K T D V Y I L N L A V A D L L L							270
360								
CTATACATC TCCCTTTG GCGTGTAT GAGTTGATG GTTGGTTT AGGCAATA AHTGCAAA TACTTCACC CTGTACACA	L F T L P F W A V N A V H G W V L G K I M C K I T S A L Y T							360
450								
CTAACTTG TCTCTGAT CAGTTTCTG GTTGTATCA GATACAG AGTGTGCA GTACTAAG TTTGACGCA ATACGGTGG	L N F V S G M Q F L A C I S I D R Y V A V T K V P S Q S G V							450
540								
GGAACACT GCTGGTAT CTTTCTGT GTCTGATG CTGCACTT CCTGACCA CCGAGTGG TTTTATAC AGTAAATAC	G K P C W I I C P C V W M A A I L L S I P Q L V F Y T V N D							540
630								
ATGCTAGGT GATTTGCT TTTGCTGG TACCTGGA CATCATGA AGCTGATT CAATCTAG AGATCTCAT TCGATTGTA	N A R C I P I F P R Y L G T S M K A L I Q M L E I C I G F V							630
720								
GTACTTTC TTTATGCG GTTGCTGC TTTATCAG CAGGACT CATGAGTG CCAACATTA AAATCTG AGCCATATA	V P F L I H G V C Y F I T A R T L M K M P N I K I S R P L K							720
810								
GTCTGCA CAGTGTAT AGTTTCAAT GTACTCAC TCCCTGTA CATTGCAAG TCTGCGAG CCAATGACT CATCTCTCC	V L L T V V I V P I V T Q L P Y N I V K P C R A I D I I Y S							810
900								
CTGATACA GCTGCAAT GAGCAACC ATGCACTG CCACTGAGT CACAGAAC ATCCACTT TTTACACTG CTTCACTCA	L I T S C N M S K R M D I A I Q V T E S I A L F H S C L N P							900
990								
ATCTTTAG TTTTATGG AGCATTTT AAACATGG TATGCAAT GGCACAAA TATCTCTT CCGACACA GAGCAAGT	I L Y V F H G A S F K N Y V M K V A K K Y G S W R R Q R Q S							990
1080								
GTGAGGAGT TTTCTTGA TTTGAGGT CACAGAGC CACAGTAC TTTGCAAT TAAGTAAA ACTCTCTCT CTTTCTTGG	V E E F P F D S E G P T E P T S T F S I							1080
1147								
GATACATG ATGTGCTT TCCCTAAA TAACTCT CCTCTCTT GAAAAAAA AAAAA								1147

3/6




FIG. 3B




FIG. 3C

4/6




FIG. 4

## STALKOKINES EXPRESSION




FIG. 5

SUBSTITUTE SHEET (RULE 26)






FIG. 6

6/6




FIG. 7

SUBSTITUTE SHEET (RULE 26)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number  
**WO 01/46698 A3**

(51) International Patent Classification<sup>7</sup>: **G01N 33/68.**  
C12N 15/10

(74) Agents: **APPLE, Randolph, T. et al.;** Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).

(21) International Application Number: **PCT/US00/34503**

(22) International Filing Date:  
18 December 2000 (18.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/172.979 20 December 1999 (20.12.1999) US  
60/186.626 3 March 2000 (03.03.2000) US  
09/721.908 24 November 2000 (24.11.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **CHEMO-CENTRYX, INC.** [US/US]; 1539 Industrial Road, San Carlos, CA 94070 (US).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SCHALL, Thomas, J.** [US/US]; 2050 Mill Ave., Menlo Park, CA 94025 (US). **TALBOT, Dale** [CA/US]; 67 Brentwood Avenue, San Francisco, CA 94127 (US). **MIAO, Zhenhua** [CN/US]; 7070 Rainbow Drive #2, San Jose, CA 95129 (US). **WEI, Zheng** [US/US]; 75 Duane Street #12, Redwood City, CA 94063 (US).

(88) Date of publication of the international search report:  
21 February 2002

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **TETHERED LIGANDS AND METHODS OF USE**




Control (-)



ELC-Stalkokine (SK)




ELC-SK + soluble ELC



ELC-SK + soluble TECK



ELC-SK + soluble SLC



ELC-SK + soluble MCP-3

(57) Abstract: The invention provides methods, reagents and devices for analysis of receptor ligand interactions, determining the profile of receptor expression in cells and cell populations, and diagnostic and drug screening methods. The invention makes use of immobilized tethered ligand fusion proteins having a ligand domain, a stalk domain, and optionally an immobilization domain.

WO 01/46698 A3

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/34503

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/68 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, BIOSIS, EPO-Internal, MEDLINE, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEN J ET AL: "TETHERED LIGAND LIBRARY FOR DISCOVERY OF PEPTIDE AGONISTS" JOURNAL OF BIOLOGICAL CHEMISTRY, US, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 270, no. 40, 6 October 1995 (1995-10-06), pages 23398-23401, XP002910821 ISSN: 0021-9258 abstract; figure 1 ---	1-23, 36-45
Y	DOORBAR, J. AND WINTER G.: "Isolation of a peptide antagonist to the thrombin receptor using phage display." J. MOL. BIOL., vol. 244, 1994, pages 361-369, XP001000952 page 363, column 2 --- -/--	1-23, 36-45



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 July 2001

Date of mailing of the international search report

19.10.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hardon, E

# INTERNATIONAL SEARCH REPORT

Inter national Application No  
PCT/US 00/34503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CHAN PO-YING ET AL: "VLA-4 integrin mediates lymphocyte migration on the inducible endothelial cell ligand VCAM-1 and the extracellular matrix ligand fibronectin." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 33, 1993, pages 24655-24664, XP000999393 ISSN: 0021-9258 the whole document</p> <p>---</p>	1-23, 36-45
Y	<p>WO 98 49557 A (ELLIS WALTHER; B E SAFE INC (US); POWERS LINDA (US)) 5 November 1998 (1998-11-05) page 15, line 15 - line 19; claim 12</p> <p>---</p>	1-23, 36-45
Y	<p>WO 97 17375 A (CHIRON DIAGNOSTICS CORP; PIRAN URI (US); CHANG STEVE C S (US); RIO) 15 May 1997 (1997-05-15) the whole document</p> <p>---</p>	1-23, 36-45
Y	<p>EP 0 626 450 A (WISCONSIN ALUMNI RES FOUND) 30 November 1994 (1994-11-30) the whole document</p> <p>---</p>	1-23, 36-45
Y	<p>WO 97 27299 A (SCHERING CORP) 31 July 1997 (1997-07-31) cited in the application the whole document</p> <p>---</p>	1-23, 36-45
Y	<p>WO 94 19694 A (JOHNSON CHARLES R; SHOEMAKER KEVIN R (US); CASS ROBERT T (US); GIE) 1 September 1994 (1994-09-01) abstract; claims 1,2</p> <p>---</p>	1-23, 36-45
Y	<p>US 5 817 527 A (RIORDAN WILLIAM J ET AL) 6 October 1998 (1998-10-06) the whole document</p> <p>---</p>	1-23, 36-45
Y	<p>BAZAN J F ET AL: "A NEW CLASS OF MEMBRANE-BOUNDED CHEMOKINE WITH A CX3C MOTIF" NATURE, GB, MACMILLAN JOURNALS LTD. LONDON, vol. 385, no. 6617, 13 February 1997 (1997-02-13), pages 640-644, XP002030092 ISSN: 0028-0836 cited in the application the whole document</p> <p>---</p>	1-23, 36-45
	<p>---</p> <p>-/--</p>	

# INTERNATIONAL SEARCH REPORT

Inter national Application No  
PCT/US 00/34503

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>GOSLING J ET AL: "Cutting edge: identification of a novel chemokine receptor that binds dendritic cell- and T cell-active chemokines including ELC, SLC, and TECK." JOURNAL OF IMMUNOLOGY, (2000 MAR 15) 164 (6) 2851-6., XP000961005 the whole document</p> <p>---</p>	1-23, 36-45
P,X	<p>DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; SCHALL, THOMAS J.: "Novel technology for the identification of specific inhibitors of orphan chemokine receptors" Database accession no. PREV2000000222786 XP002171801 abstract &amp; ABSTRACTS OF PAPERS AMERICAN CHEMICAL SOCIETY, vol. 219, no. 1-2, 26 - 30 March 2000, pages MEDI-339,</p> <p>-----</p>	1-23, 36-45

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/34503

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-23, 36-45

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-5, 36-45

assay device comprising a plurality of different immobilized "tethered" (attached) ligand fusion proteins organized in an array, wherein the ligand fusion protein contains a "stalk" (spacer), and related matter (independent claims 1 and 36; and dependent claims), and tethered ligand fusion protein comprising a stalk (independent claim 44, and dependent claim 45).

2. Claims: 6-23

method for identifying an interaction between a receptor and a ligand comprising contacting a cell expressing a receptor with an immobilized "tethered" ligand fusion protein (no stalk, no array), and related matter (independent claims 6, 13 and 16; and dependent claims).

3. Claims: 24-35

method for diagnosis, and method for detecting the effect of a drug, said method comprising "determining a receptor profile" (independent claims 24 and 31; and dependent claims),



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/34503

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9849557	A	05-11-1998	AU 7164998 A WO 9849557 A1	24-11-1998 05-11-1998
WO 9717375	A	15-05-1997	US 5817527 A AU 7327796 A EP 0859793 A1 WO 9717375 A1 JP 2000500441 T	06-10-1998 29-05-1997 26-08-1998 15-05-1997 18-01-2000
EP 0626450	A	30-11-1994	EP 0626450 A2 JP 7147987 A US 5851791 A	30-11-1994 13-06-1995 22-12-1998
WO 9727299	A	31-07-1997	AU 1825097 A EP 0876487 A1 JP 11503030 T WO 9727299 A1	20-08-1997 11-11-1998 23-03-1999 31-07-1997
WO 9419694	A	01-09-1994	US 5576220 A US 5585275 A AU 6393994 A JP 8507602 T WO 9419694 A1 AU 4844593 A WO 9405394 A1	19-11-1996 17-12-1996 14-09-1994 13-08-1996 01-09-1994 29-03-1994 17-03-1994
US 5817527	A	06-10-1998	AU 7327796 A EP 0859793 A1 WO 9717375 A1 JP 2000500441 T	29-05-1997 26-08-1998 15-05-1997 18-01-2000

Form PCT/ISA/210 (patent family annex) (July 1992)